

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

International Patent Classification 6 :

C12N 15/00

A2

(11) International Publication Number:

WO 99/29837

(43) International Publication Date:

17 June 1999 (17.06.99)

(21) International Application Number:

PCT/EP98/07945

(22) International Filing Date:

5 December 1998 (07.12.98)

(30) Priority Data:

97121462.2

5 December 1997 (05.12.97)

EP

98118756.0

5 October 1998 (05.10.98)

EP

(71) Applicant (for all designated States except US): EUROPAISCHES LABORATORIUM FÜR MOLEKULARBIOLOGIE (EMBL) [DE/DE]; Meyerhofstrasse 1, D-69117 Heidelberg (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): STEWART, Francis [AU/DE]; Lärchenweg 3, D-69181 Leimen (DE). ZHANG, Youming [CN/DE]; Friedrich-Ebert-Anlage 51e, D-69117 Heidelberg (DE). BUCHHOLZ, Frank [DE/DE]; Neuenkirchener Weg 44a, D-28779 Bremen (DE).

(74) Agents: WEICKMANN, H. et al.; Kopernikusstrasse 9, D-81679 München (DE).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW. ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: NOVEL DNA CLONING METHOD

(57) Abstract

The invention refers to a novel method for cloning DNA molecules using a homologous recombination mechanism between at least two DNA molecules comprising: a) providing a host cell capable of performing homologous recombination, b) contacting in said host cell a first DNA molecule which is capable of being replicated in said host cell with a second DNA molecule comprising at least two regions of sequence homology to regions on the first DNA molecule, under conditions which favour homologous recombination between said first and second DNA molecules and c) selecting a host cell in which homologous recombination between said first and second DNA molecules has occurred.

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



- 5 -

In the method of the present invention the homologous recombination preferably occurs via the *recET* mechanism, i.e. the homologous recombination is mediated by the gene products of the *recE* and the *recT* genes which are preferably selected from the *E. coli* genes *recE* and *recT* or functionally related genes such as the phage  $\lambda$  *red $\alpha$*  and *red $\beta$*  genes.

The host cell suitable for the method of the present invention preferably is a bacterial cell, e.g. a gram-negative bacterial cell. More preferably, the host cell is an enterobacterial cell, such as *Salmonella*, *Klebsiella* or *Escherichia*. Most preferably the host cell is an *Escherichia coli* cell. It should be noted, however, that the cloning method of the present invention is also suitable for eukaryotic cells, such as fungi, plant or animal cells.

Preferably, the host cell used for homologous recombination and propagation of the cloned DNA can be any cell, e.g. a bacterial strain in which the products of the *recE* and *recT*, or *red $\alpha$*  and *red $\beta$* , genes are expressed. The host cell may comprise the *recE* and *recT* genes located on the host cell chromosome or on non-chromosomal DNA, preferably on a vector, e.g. a plasmid. In a preferred case, the *RecE* and *RecT*, or *Red $\alpha$*  and *Red $\beta$* , gene products are expressed from two different regulatable promoters, such as the arabinose-inducible *BAD* promoter or the *lac* promoter or from non-regulatable promoters. Alternatively, the *recE* and *recT*, or *red $\alpha$*  and *red $\beta$* , genes are expressed on a polycistronic mRNA from a single regulatable or non-regulatable promoter. Preferably the expression is controlled by regulatable promoters.

Especially preferred is also an embodiment, wherein the *recE* or *red $\alpha$*  gene is expressed by a regulatable promoter. Thus, the recombinogenic potential of the system is only elicited when required and, at other times, possible undesired recombination reactions are limited. The *recT* or *red $\beta$*  gene, on the other hand, is preferably overexpressed with respect to *recE* or *red $\alpha$* . This may be accomplished by using a strong constitutive promoter, e.g. the

EM7 promoter and/or by using a higher copy number of *recT*, or *red $\beta$* , versus *recE*, or *red $\alpha$* , genes.

For the purpose of the present invention any *recE* and *recT* genes are  
5 suitable insofar as they allow a homologous recombination of first and  
second DNA molecules with sufficient efficiency to give rise to  
recombination products in more than 1 in 10<sup>9</sup> cells transfected with DNA.  
The *recE* and *recT* genes may be derived from any bacterial strain or from  
bacteriophages or may be mutants and variants thereof. Preferred are *recE*  
10 and *recT* genes which are derived from *E.coli* or from *E.coli* bacteriophages,  
such as the *red $\alpha$*  and *red $\beta$*  genes from lambdoid phages, e.g. bacteriophage  
 $\lambda$ .

More preferably, the *recE* or *red $\alpha$*  gene is selected from a nucleic acid  
15 molecule comprising

- (a) the nucleic acid sequence from position 1320 (ATG) to 2159 (GAC) as depicted in Fig.7B,
- (b) the nucleic acid sequence from position 1320 (ATG) to 1998(CGA) as depicted in Fig.14B,
- 20 (c) a nucleic acid encoding the same polypeptide within the degeneracy of the genetic code and/or
- (d) a nucleic acid sequence which hybridizes under stringent conditions with the nucleic acid sequence from (a), (b) and/or (c).

25 More preferably, the *recT* or *red $\beta$*  gene is selected from a nucleic acid molecule comprising

- (a) the nucleic acid sequence from position 2155 (ATG) to 2961 (GAA) as depicted in Fig.7B,
- (b) the nucleic acid sequence from position 2086 (ATG) to 2868 (GCA) as  
30 depicted in Fig.14B,
- (c) a nucleic acid encoding the same polypeptide within the degeneracy of the genetic code and/or

- 7 -

(d) a nucleic acid sequence which hybridizes under stringent conditions with the nucleic acid sequences from (a), (b) and/or (c).

It should be noted that the present invention also encompasses mutants and variants of the given sequences, e.g. naturally occurring mutants and variants or mutants and variants obtained by genetic engineering. Further it should be noted that the *recE* gene depicted in Fig.7B is an already truncated gene encoding amino acids 588-866 of the native protein. Mutants and variants preferably have a nucleotide sequence identity of at least 60%, preferably of at least 70% and more preferably of at least 80% of the *recE* and *recT* sequences depicted in Fig.7B and 13B, and of the *red $\alpha$*  and *red $\beta$*  sequences depicted in Fig.14B.

According to the present invention hybridization under stringent conditions preferably is defined according to Sambrook et al. (1989), *infra*, and comprises a detectable hybridization signal after washing for 30 min in 0.1 x SSC, 0.5% SDS at 55°C, preferably at 62°C and more preferably at 63°C.

In a preferred case the *recE* and *recT* genes are derived from the corresponding endogenous genes present in the *E.coli* K12 strain and its derivatives or from bacteriophages. In particular, strains that carry the *sbcA* mutation are suitable. Examples of such strains are JC8679 and JC 9604 (Gillen et al. (1981), *supra*). Alternatively, the corresponding genes may also be obtained from other coliphages such as lambdoid phages or phage P22.

The genotype of JC 8679 and JC 9604 is Sex (Hfr, F+, F-, or F') : F-. JC 8679 comprises the mutations: *recBC* 21, *recC* 22, *sbcA* 23, *thr*-1, *ara*-14, *leu* B 6, *DE* (*gpt-proA*) 62, *lacY*1, *tsx*-33, *gluV*44 (AS), *galK*2 (Oc), LAM-, *his*-60, *relA* 1, *rps* L31 (*strR*), *xyl* A5, *mtl*-1, *argE*3 (Oc) and *thi*-1. JC 9604 comprises the same mutations and further the mutation *recA* 56.

Further, it should be noted that the *recE* and *recT*, or *red $\alpha$*  and *red $\beta$* , genes can be isolated from a first donor source, e.g. a donor bacterial cell and transformed into a second receptor source, e.g. a receptor bacterial or eukaryotic cell in which they are expressed by recombinant DNA means.

In one embodiment of the invention, the host cell used is a bacterial strain having an *sbcA* mutation, e.g. one of *E.coli* strains JC 8679 and JC 9604 mentioned above. However, the method of the invention is not limited to host cells having an *sbcA* mutation or analogous cells. Surprisingly, it has been found that the cloning method of the invention also works in cells without *sbcA* mutation, whether *recBC* + or *recBC* -, e.g. also in prokaryotic *recBC* + host cells, e.g. in *E.coli* *recBC* + cells. In that case preferably those host cells are used in which the product of a *recBC* type exonuclease inhibitor gene is expressed. Preferably, the exonuclease inhibitor is capable of inhibiting the host *recBC* system or an equivalent thereof. A suitable example of such exonuclease inhibitor gene is the  $\lambda$  *red $\gamma$*  gene (Murphy, J.Bacteriol. 173 (1991), 5808-5821) and functional equivalents thereof, respectively, which, for example, can be obtained from other coliphages such as from phage P22 (Murphy, J.Biol.Chem.269 (1994), 22507-22516).

More preferably, the exonuclease inhibitor gene is selected from a nucleic acid molecule comprising

(a) the nucleic acid sequence from position 3588 (ATG) to 4002 (GTA) as depicted in Fig. 14A,

(b) a nucleic acid encoding the same polypeptide within the degeneracy of the genetic code and/or

(c) a nucleic acid sequence which hybridizes under stringent conditions (as defined above) with the nucleic acid sequence from (a) and/ or (b).

Surprisingly, it has been found that the expression of an exonuclease inhibitor gene in both *recBC* + and *recBC* - strains leads to significant improvement of cloning efficiency.

- 9 -

The cloning method according to the present invention employs a homologous recombination between a first DNA molecule and a second DNA molecule. The first DNA molecule can be any DNA molecule that carries an origin of replication which is operative in the host cell, e.g. an E.coli replication origin. Further, the first DNA molecule is present in a form which is capable of being replicated in the host cell. The first DNA molecule, i.e. the vector, can be any extrachromosomal DNA molecule containing an origin of replication which is operative in said host cell, e.g. a plasmid including single, low, medium or high copy plasmids or other extrachromosomal circular DNA molecules based on cosmid, P1, BAC or PAC vector technology. Examples of such vectors are described, for example, by Sambrook et al. (Molecular Cloning, Laboratory Manual, 2nd Edition (1989), Cold Spring Harbor Laboratory Press) and Ioannou et al. (Nature Genet. 6 (1994), 84-89) or references cited therein. The first DNA molecule can also be a host cell chromosome, particularly the E.coli chromosome. Preferably, the first DNA molecule is a double-stranded DNA molecule.

The second DNA molecule is preferably a linear DNA molecule and comprises at least two regions of sequence homology, preferably of sequence identity to regions on the first DNA molecule. These homology or identity regions are preferably at least 15 nucleotides each, more preferably at least 20 nucleotides and, most preferably, at least 30 nucleotides each. Especially good results were obtained when using sequence homology regions having a length of about 40 or more nucleotides, e.g. 60 or more nucleotides. The two sequence homology regions can be located on the linear DNA fragment so that one is at one end and the other is at the other end, however they may also be located internally. Preferably, also the second DNA molecule is a double-stranded DNA molecule.

30

The two sequence homology regions are chosen according to the experimental design. There are no limitations on which regions of the first

- 10 -

DNA molecule can be chosen for the two sequence homology regions located on the second DNA molecule, except that the homologous recombination event cannot delete the origin of replication of the first DNA molecule. The sequence homology regions can be interrupted by non-identical sequence regions as long as sufficient sequence homology is retained for the homologous recombination reaction. By using sequence homology arms having non-identical sequence regions compared to the target site mutations such as substitutions, e.g. point mutations, insertions and/or deletions may be introduced into the target site by ET cloning.

10

The second foreign DNA molecule which is to be cloned in the bacterial cell may be derived from any source. For example, the second DNA molecule may be synthesized by a nucleic acid amplification reaction such as a PCR where both of the DNA oligonucleotides used to prime the amplification contain in addition to sequences at the 3'-ends that serve as a primer for the amplification, one or the other of the two homology regions. Using oligonucleotides of this design, the DNA product of the amplification can be any DNA sequence suitable for amplification and will additionally have a sequence homology region at each end.

20

A specific example of the generation of the second DNA molecule is the amplification of a gene that serves to convey a phenotypic difference to the bacterial host cells, in particular, antibiotic resistance. A simple variation of this procedure involves the use of oligonucleotides that include other sequences in addition to the PCR primer sequence and the sequence homology region. A further simple variation is the use of more than two amplification primers to generate the amplification product. A further simple variation is the use of more than one amplification reaction to generate the amplification product. A further variation is the use of DNA fragments obtained by methods other than PCR, for example, by endonuclease or restriction enzyme cleavage to linearize fragments from any source of DNA.

30

- 11 -

It should be noted that the second DNA molecule is not necessarily a single species of DNA molecule. It is of course possible to use a heterogeneous population of second DNA molecules, e.g. to generate a DNA library, such as a genomic or cDNA library.

5

The method of the present invention may comprise the contacting of the first and second DNA molecules in vivo. In one embodiment of the present invention the second DNA fragment is transformed into a bacterial strain that already harbors the first vector DNA molecule. In a different  
10 embodiment, the second DNA molecule and the first DNA molecule are mixed together in vitro before co-transformation in the bacterial host cell. These two embodiments of the present invention are schematically depicted in Fig. 1. The method of transformation can be any method known in the art (e.g. Sambrook et al. supra). The preferred method of transformation or co-  
15 transformation, however, is electroporation.

After contacting the first and second DNA molecules under conditions which favour homologous recombination between first and second DNA molecules via the ET cloning mechanism a host cell is selected, in which  
20 homologous recombination between said first and second DNA molecules has occurred. This selection procedure can be carried out by several different methods. In the following three preferred selection methods are depicted in Fig.2 and described in detail below.

25 In a first selection method a second DNA fragment is employed which carries a gene for a marker placed between the two regions of sequence homology wherein homologous recombination is detectable by expression of the marker gene. The marker gene may be a gene for a phenotypic marker which is not expressed in the host or from the first DNA molecule.  
30 Upon recombination by ET cloning, the change in phenotype of the host strain conveyed by the stable acquisition of the second DNA fragment identifies the ET cloning product.

In a preferred case, the phenotypic marker is a gene that conveys resistance to an antibiotic, in particular, genes that convey resistance to kanamycin, ampicillin, chloramphenicol, tetracyclin or any other substance that shows bacteriocidal or bacteriostatic effects on the bacterial strain employed.

5

A simple variation is the use of a gene that complements a deficiency present within the bacterial host strain employed. For example, the host strain may be mutated so that it is incapable of growth without a metabolic supplement. In the absence of this supplement, a gene on the second DNA  
10 fragment can complement the mutational defect thus permitting growth. Only those cells which contain the episome carrying the intended DNA rearrangement caused by the ET cloning step will grow.

In another example, the host strain carries a phenotypic marker gene which  
15 is mutated so that one of its codons is a stop codon that truncates the open reading frame. Expression of the full length protein from this phenotypic marker gene requires the introduction of a suppressor tRNA gene which, once expressed, recognizes the stop codon and permits translation of the full open reading frame. The suppressor tRNA gene is introduced by the ET  
20 cloning step and successful recombinants identified by selection for, or identification of, the expression of the phenotypic marker gene. In these cases, only those cells which contain the intended DNA rearrangement caused by the ET cloning step will grow.

25 A further simple variation is the use of a reporter gene that conveys a readily detectable change in colony colour or morphology. In a preferred case, the green fluorescence protein (GFP) can be used and colonies carrying the ET cloning product identified by the fluorescence emissions of GFP. In another preferred case, the lacZ gene can be used and colonies  
30 carrying the ET cloning product identified by a blue colony colour when X-gal is added to the culture medium.



- 13 -

In a second selection method the insertion of the second DNA fragment into the first DNA molecule by ET cloning alters the expression of a marker present on the first DNA molecule. In this embodiment the first DNA molecule contains at least one marker gene between the two regions of sequence homology and homologous recombination may be detected by an altered expression, e.g. lack of expression of the marker gene.

In a preferred application, the marker present on the first DNA molecule is a counter-selectable gene product, such as the *sacB*, *ccdB* or tetracycline-resistance genes. In these cases, bacterial cells that carry the first DNA molecule unmodified by the ET cloning step after transformation with the second DNA fragment, or co-transformation with the second DNA fragment and the first DNA molecule, are plated onto a medium so the expression of the counter-selectable marker conveys a toxic or bacteriostatic effect on the host. Only those bacterial cells which contain the first DNA molecule carrying the intended DNA rearrangement caused by the ET cloning step will grow.

In another preferred application, the first DNA molecule carries a reporter gene that conveys a readily detectable change in colony colour or morphology. In a preferred case, the green fluorescence protein (GFP) can be present on the first DNA molecule and colonies carrying the first DNA molecule with or without the ET cloning product can be distinguished by differences in the fluorescence emissions of GFP. In another preferred case, the *lacZ* gene can be present on the first DNA molecule and colonies carrying the first DNA molecule with or without the ET cloning product identified by a blue or white colony colour when X-gal is added to the culture medium.

In a third selection method the integration of the second DNA fragment into the first DNA molecule by ET cloning removes a target site for a site specific recombinase, termed here an RT (for recombinase target) present

on the first DNA molecule between the two regions of sequence homology. A homologous recombination event may be detected by removal of the target site.

5 In the absence of the ET cloning product, the RT is available for use by the corresponding site specific recombinase. The difference between the presence or not of this RT is the basis for selection of the ET cloning product. In the presence of this RT and the corresponding site specific recombinase, the site specific recombinase mediates recombination at this  
10 RT and changes the phenotype of the host so that it is either not able to grow or presents a readily observable phenotype. In the absence of this RT, the corresponding site specific recombinase is not able to mediate recombination.

15 In a preferred case, the first DNA molecule to which the second DNA fragment is directed, contains two RTs, one of which is adjacent to, but not part of, an antibiotic resistance gene. The second DNA fragment is directed, by design, to remove this RT. Upon exposure to the corresponding site specific recombinase, those first DNA molecules that do not carry the ET  
20 cloning product will be subject to a site specific recombination reaction between the RTs that remove the antibiotic resistance gene and therefore the first DNA molecule fails to convey resistance to the corresponding antibiotic. Only those first DNA molecules that contain the ET cloning product, or have failed to be site specifically recombined for some other  
25 reason, will convey resistance to the antibiotic.

In another preferred case, the RT to be removed by ET cloning of the second DNA fragment is adjacent to a gene that complements a deficiency present within the host strain employed. In another preferred case, the RT  
30 to be removed by ET cloning of the second DNA fragment is adjacent to a reporter gene that conveys a readily detectable change in colony colour or morphology.

- 15 -

In another preferred case, the RT to be removed by ET cloning of the second DNA fragment is anywhere on a first episomal DNA molecule and the episome carries an origin of replication incompatible with survival of the bacterial host cell if it is integrated into the host genome. In this case the host genome carries a second RT, which may or may not be a mutated RT so that the corresponding site specific recombinase can integrate the episome, via its RT, into the RT sited in the host genome. Other preferred RTs include RTs for site specific recombinases of the resolvase/transposase class. RTs include those described from existing examples of site specific recombination as well as natural or mutated variations thereof.

The preferred site specific recombinases include Cre, FLP, Kw or any site specific recombinase of the integrase class. Other preferred site specific recombinases include site specific recombinases of the resolvase/transposase class.

There are no limitations on the method of expression of the site specific recombinase in the host cell. In a preferred method, the expression of the site specific recombinase is regulated so that expression can be induced and quenched according to the optimisation of the ET cloning efficiency. In this case, the site specific recombinase gene can be either integrated into the host genome or carried on an episome. In another preferred case, the site specific recombinase is expressed from an episome that carries a conditional origin of replication so that it can be eliminated from the host cell.

In another preferred case, at least two of the above three selection methods are combined. A particularly preferred case involves a two-step use of the first selection method above, followed by use of the second selection method. This combined use requires, most simply, that the DNA fragment to be cloned includes a gene, or genes that permits the identification, in the first step, of correct ET cloning products by the acquisition of a phenotypic

change. In a second step, expression of the gene or genes introduced in the first step is altered so that a second round of ET cloning products can be identified. In a preferred example, the gene employed is the tetracycline resistance gene and the first step ET cloning products are identified by the acquisition of tetracycline resistance. In the second step, loss of expression of the tetracycline gene is identified by loss of sensitivity to nickel chloride, fusaric acid or any other agent that is toxic to the host cell when the tetracycline gene is expressed. This two-step procedure permits the identification of ET cloning products by first the integration of a gene that conveys a phenotypic change on the host, and second by the loss of a related phenotypic change, most simply by removal of some of the DNA sequences integrated in the first step. Thereby the genes used to identify ET cloning products can be inserted and then removed to leave ET cloning products that are free of these genes.

In a further embodiment of the present invention the ET cloning may also be used for a recombination method comprising the steps of

- a) providing a source of RecE and RecT, or Red $\alpha$  and Red $\beta$ , proteins,
- b) contacting a first DNA molecule which is capable of being replicated in a suitable host cell with a second DNA molecule comprising at least two regions of sequence homology to regions on the first DNA molecule, under conditions which favour homologous recombination between said first and second DNA molecules and
- c) selecting DNA molecules in which a homologous recombination between said first and second DNA molecules has occurred.

The source of RecE and RecT, or Red $\alpha$  and Red $\beta$ , proteins may be either purified or partially purified RecE and RecT, or Red $\alpha$  and Red $\beta$ , proteins or cell extracts comprising RecE and RecT, or Red $\alpha$  and Red $\beta$ , proteins.

The homologous recombination event in this embodiment may occur in vitro, e.g. when providing a cell extract containing further components

- 17 -

required for homologous recombination. The homologous recombination event, however, may also occur in vivo, e.g. by introducing RecE and RecT, or Red $\alpha$  and Red $\beta$ , proteins or the extract in a host cell (which may be recET positive or not, or red $\alpha\beta$  positive or not) and contacting the DNA molecules in the host cell. When the recombination occurs in vitro the selection of DNA molecules may be accomplished by transforming the recombination mixture in a suitable host cell and selecting for positive clones as described above. When the recombination occurs in vivo the selection methods as described above may directly be applied.

A further subject matter of the invention is the use of cells, preferably bacterial cells, most preferably, E.coli cells capable of expressing the recE and recT, or red $\alpha$  and red $\beta$ , genes as a host cell for a cloning method involving homologous recombination.

Still a further subject matter of the invention is a vector system capable of expressing recE and recT, or red $\alpha$  and red $\beta$ , genes in a host cell and its use for a cloning method involving homologous recombination. Preferably, the vector system is also capable of expressing an exonuclease inhibitor gene as defined above, e.g. the  $\lambda$  red $\gamma$  gene. The vector system may comprise at least one vector. The recE and recT, or red $\alpha$  and red $\beta$ , genes are preferably located on a single vector and more preferably under control of a regulatable promoter which may be the same for both genes or a single promoter for each gene. Especially preferred is a vector system which is capable of overexpressing the recT, or red $\beta$ , gene versus the recE, or red $\alpha$ , gene.

Still a further subject matter of the invention is the use of a source of RecE and RecT, or Red $\alpha$  and Red $\beta$ , proteins for a cloning method involving homologous recombination.

A still further subject matter of the invention is a reagent kit for cloning comprising

- (a) a host cell, preferably a bacterial host cell,
- (b) means of expressing *recE* and *recT*, or *red $\alpha$*  and *red $\beta$* , genes in said host cell, e.g. comprising a vector system, and
- (c) a recipient cloning vehicle, e.g. a vector, capable of being replicated in said cell.

On the one hand, the recipient cloning vehicle which corresponds to the first DNA molecule of the process of the invention can already be present in the bacterial cell. On the other hand, it can be present separated from the bacterial cell.

In a further embodiment the reagent kit comprises

- (a) a source for *RecE* and *RecT*, or *Red $\alpha$*  and *Red $\beta$* , proteins and
- (b) a recipient cloning vehicle capable of being propagated in a host cell and
- (c) optionally a host cell suitable for propagating said recipient cloning vehicle.

The reagent kit furthermore contains, preferably, means for expressing a site specific recombinase in said host cell, in particular, when the recipient ET cloning product contains at least one site specific recombinase target site. Moreover, the reagent kit can also contain DNA molecules suitable for use as a source of linear DNA fragments used for ET cloning, preferably by serving as templates for PCR generation of the linear fragment, also as specifically designed DNA vectors from which the linear DNA fragment is released by restriction enzyme cleavage, or as prepared linear fragments included in the kit for use as positive controls or other tasks. Moreover, the reagent kit can also contain nucleic acid amplification primers comprising a region of homology to said vector. Preferably, this region of homology is located at the 5'-end of the nucleic acid amplification primer.

- 19 -

The invention is further illustrated by the following Sequence listings, Figures and Examples.

- SEQ ID NO. 1: shows the nucleic acid sequence of the plasmid pBAD24-rec ET (Fig. 7).
- 5 SEQ ID NOs 2/3: show the nucleic acid and amino acid sequences of the truncated recE gene (t-recE) present on pBAD24-recET at positions 1320-2162.
- SEQ ID NOs 4/5: show the nucleic acid and amino acid sequences of the recT gene present on pBAD24-recET at position 2155-2972.
- 10 SEQ ID NOs 6/7: show the nucleic acid and amino acid sequences of the araC gene present on the complementary stand to the one shown of pBAD24-recET at positions 974-996.
- SEQ ID NOs 8/9: show the nucleic acid an amino acid sequences of the bla gene present on pBAD24-recET at positions 3493-4353.
- 15 SEQ ID NO 10: shows the nucleic acid sequence of the plasmid pBAD-ET $\gamma$  (Fig. 13).
- SEQ ID No 11: shows the nucleic acid sequence of the plasmid pBAD- $\alpha\beta\gamma$  (Fig. 14) as well as the coding regions for the genes red $\alpha$  (1320-200), red $\beta$  (2086-2871) and red $\gamma$  (3403-3819).
- 20 SEQ ID NOs 12-14: show the amino acid sequences of the Red $\alpha$ , Red $\beta$  and Red $\gamma$  proteins, respectively. The red $\gamma$  sequence is present on each of pBAD-ET $\gamma$  (Fig. 13) and pBAD- $\alpha\beta\gamma$  (Fig. 14).
- 25

### Figure 1

A preferred method for ET cloning is shown by diagram. The linear DNA  
30 fragment to be cloned is synthesized by PCR using oligonucleotide primers that contain a left homology arm chosen to match sequences in the recipient episome and a sequence for priming in the PCR reaction, and a

- 20 -

right homology arm chosen to match another sequence in the recipient episome and a sequence for priming in the PCR reaction. The product of the PCR reaction, here a selectable marker gene (sm1), is consequently flanked by the left and right homology arms and can be mixed together in vitro with the episome before co-transformation, or transformed into a host cell harboring the target episome. The host cell contains the products of the *recE* and *recT* genes. ET cloning products are identified by the combination of two selectable markers, sm1 and sm2 on the recipient episome.

## Figure 2

Three ways to identify ET cloning products are depicted. The first, (on the left of the figure), shows the acquisition, by ET cloning, of a gene that conveys a phenotypic difference to the host, here a selectable marker gene (sm). The second (in the centre of the figure) shows the loss, by ET cloning, of a gene that conveys a phenotypic difference to the host, here a counter selectable marker gene (counter-sm). The third shows the loss of a target site (RT, shown as triangles on the circular episome) for a site specific recombinase (SSR), by ET cloning. In this case, the correct ET cloning product deletes one of the target sites required by the SSR to delete a selectable marker gene (sm). The failure of the SSR to delete the sm gene identifies the correct ET cloning product.

## Figure 3

A simple example of ET cloning is presented.

(a) Top panel - PCR products (left lane) synthesized from oligonucleotides designed as described in Fig.1 to amplify by PCR a kanamycin resistance gene and to be flanked by homology arms present in the recipient vector, were mixed in vitro with the recipient vector (2nd lane) and cotransformed into a *recET+* *E.coli* host. The recipient vector carried an ampicillin



- 21 -

resistance gene. (b) Transformation of the *sbca* E.coli strain JC9604 with either the PCR product alone (0.2  $\mu$ g) or the vector alone (0.3  $\mu$ g) did not convey resistance to double selection with ampicillin and kanamycin (amp + kan), however cotransformation of both the PCR product and the vector produced double resistant colonies. More than 95% of these colonies contained the correct ET cloning product where the kanamycin gene had precisely integrated into the recipient vector according to the choice of homology arms. The two lanes on the right of (a) show Pvu II restriction enzyme digestion of the recipient vector before and after ET cloning. (c) As for b, except that six PCR products (0.2  $\mu$ g each) were cotransformed with pSVpaZ11 (0.3  $\mu$ g each) into JC9604 and plated onto Amp + Kan plates or Amp plates. Results are plotted as Amp + Kan-resistant colonies, representing recombination products, divided by Amp-resistant colonies, representing the plasmid transformation efficiency of the competent cell preparation,  $\times 10^5$ . The PCR products were equivalent to the a-b PCR product except that homology arm lengths were varied. Results are from five experiments that used the same batches of competent cells and DNAs. Error bars represent standard deviation. (d) Eight products flanked by 50 bp homology arms were cotransformed with pSVpaZ11 into JC9604. All eight PCR products contained the same left homology arm and amplified neo gene. The right homology arms were chosen from the pSVpaZ11 sequence to be adjacent to (0), or at increasing distances (7-3100 bp), from the left. Results are from four experiments.

Figure 4

ET cloning in an approximately 100kb P1 vector to exchange the selectable marker.

A P1 clone which uses a kanamycin resistance gene as selectable marker and which contains at least 70kb of the mouse Hox a gene cluster was used. Before ET cloning, this episome conveys kanamycin resistance (top

panel, upper left) to its host E.coli which are ampicillin sensitive (top panel, upper right). A linear DNA fragment designed to replace the kanamycin resistance gene with an ampicillin resistance gene was made by PCR as outlined in Fig.1 and transformed into E.coli host cells in which the recipient Hox a/P1 vector was resident. ET cloning resulted in the deletion of the kanamycin resistance gene, and restoration of kanamycin sensitivity (top panel, lower left) and the acquisition of ampicillin resistance (top panel, lower right). Precise DNA recombination was verified by restriction digestion and Southern blotting analyses of isolated DNA before and after ET cloning (lower panel).

#### Figure 5

ET cloning to remove a counter selectable marker

A PCR fragment (upper panel, left, third lane) made as outlined in Figs.1 and 2 to contain the kanamycin resistance gene was directed by its chosen homology arms to delete the counter selectable *ccdB* gene present in the vector, pZero-2.1. The PCR product and the pZero vector were mixed in vitro (upper panel, left, 1st lane) before cotransformation into a *recE/recT* + E.coli host. Transformation of pZero-2.1 alone and plating onto kanamycin selection medium resulted in little colony growth (lower panel, left). Cotransformation of pZero-2.1 and the PCR product presented ET cloning products (lower panel, right) which showed the intended molecular event as visualized by *Pvu* II digestion (upper panel, right).

#### Figure 6

ET cloning mediated by inducible expression of *recE* and *recT* from an episome.

*RecE/RecT* mediate homologous recombination between linear and circular DNA molecules. (a) The plasmid pBAD24-*recET* was transformed into E.coli JC5547, and then batches of competent cells were prepared after induction of *RecE/RecT* expression by addition of L-arabinose for the times indicated

- 23 -

before harvesting. A PCR product, made using oligonucleotides e and f to contain the chloramphenicol resistance gene (cm) of pMAK705 and 50 bp homology arms chosen to flank the ampicillin resistance gene (bla) of pBAD24-recET, was then transformed and recombinants identified on chloramphenicol plates. (b) Arabinose was added to cultures of pBAD24-recET transformed JC5547 for different times immediately before harvesting for competent cell preparation. Total protein expression was analyzed by SDS-PAGE and Coomassie blue staining. (c) The number of chloramphenicol resistant colonies per  $\mu\text{g}$  of PCR product was normalized against a control for transformation efficiency, determined by including 5 pg pZero2.1, conveying kanamycin resistance, in the transformation and plating an aliquot onto Kan plates.

#### Figure 7A

The plasmid pBAD24-recET is shown by diagram. The plasmid contains the genes recE (in a truncated form) and recT under control of the inducible BAD promoter ( $P_{BAD}$ ). The plasmid further contains an ampicillin resistance gene (Amp<sup>r</sup>) and an araC gene.

#### Figure 7B

The nucleic acid sequence and the protein coding portions of pBAD24-recET are depicted.

#### Figure 8

Manipulation of a large E.coli episome by multiple recombination steps. a Scheme of the recombination reactions. A P1 clone of the Mouse Hoxa complex, resident in JC9604, was modified by recombination with PCR products that contained the neo gene and two Flp recombination targets

(FRTs). The two PCR products were identical except that one was flanked by g and h homology arms (insertion), and the other was flanked by i and h homology arms (deletion). In a second step, the neo gene was removed by Flp recombination between the FRTs by transient transformation of a Flp expression plasmid based on the pSC101 temperature-sensitive origin (ts ori). b Upper panel; ethidium bromide stained agarose gel showing EcoR1 digestions of P1 DNA preparations from three independent colonies for each step. Middle panel; a Southern blot of the upper panel hybridized with a neo gene probe. Lower panel; a Southern blot of the upper panel hybridized with a Hoxa3 probe to visualize the site of recombination. Lanes 1, the original Hoxa3 P1 clone grown in E.coli strain NS3145. Lanes 2, replacement of the Tn903 kanamycin resistance gene resident in the P1 vector with an ampicillin resistance gene increased the 8.1 kb band (lanes 1), to 9.0 kb. Lanes 3, insertion of the Tn5-neo gene with g-h homology arms upstream of Hoxa3, increased the 6.7 kb band (lanes 1,2) to 9.0 kb. Lanes 4, Flp recombinase deleted the g-h neo gene reducing the 9.0 kb band (lanes 3) back to 6.7 kb. Lanes 5, deletion of 6 kb of Hoxa3 - 4 intergenic DNA by replacement with the i-h neo gene, decreased the 6.7 kb band (lanes 2) to 4.5 kb. Lanes 6, Flp recombinase deleted the i-h neo gene reducing the 4.5 kb band to 2.3 kb.

### Figure 9

Manipulation of the E.coli chromosome. A Scheme of the recombination reactions. The endogenous lacZ gene of JC9604 at 7.8' of the E.coli chromosome, shown in expanded form with relevant Ava I sites and coordinates, was targeted by a PCR fragment that contained the neo gene flanked by homology arms j and k, and loxP sites, as depicted. Integration of the neo gene removed most of the lacZ gene including an Ava I site to alter the 1443 and 3027 bp bands into a 3277 bp band. In a second step, the neo gene was removed by Cre recombination between the loxPs by transient transformation of a Cre expression plasmid based on the pSC101

- 25 -

temperature-sensitive origin (ts ori). Removal of the neo gene by Cre recombinase reduces the 3277 band to 2111 bp. b  $\beta$ -galactosidase expression evaluated by streaking colonies on X-Gal plates. The top row of three streaks show  $\beta$ -galactosidase expression in the host JC9604 strain (w.t.), the lower three rows (Km) show 24 independent primary colonies, 20 of which display a loss of  $\beta$ -galactosidase expression indicative of the intended recombination event. c Southern analysis of E.coli chromosomal DNA digested with Ava I using a random primed probe made from the entire lacZ coding region; lanes 1,2, w.t.; lanes 3-6, four independent white colonies after integration of the j-k neo gene; lanes 7-10; the same four colonies after transient transformation with the Cre expression plasmid.

Figure 10

Two rounds of ET cloning to introduce a point mutation. a Scheme of the recombination reactions. The lacZ gene of pSVpaX1 was disrupted in JC9604lacZ, a strain made by the experiment of Fig.9 to ablate endogenous lacZ expression and remove competitive sequences, by a sacB-neo gene cassette, synthesized by PCR to pIB279 and flanked by l and m' homology arms. The recombinants, termed pSV-sacB-neo, were selected on Amp + Kan plates. The lacZ gene of pSV-sacB-neo was then repaired by a PCR fragment made from the intact lacZ gene using l' and m' homology arms. The m' homology arm included a silent C to G change that created a BamH1 site. The recombinants, termed pSVpaX1', were identified by counter selection against the sacB gene using 7% sucrose. b  $\beta$ -galactosidase expression from pSVpaX1 was disrupted in pSV-sacB-neo and restored in pSVpaX1'. Expression was analyzed on X-gal plates. Three independent colonies of each pSV-sacB-neo and pSVpaX1' are shown. c Ethidium bromide stained agarose gels of BamH1 digested DNA prepared from independent colonies taken after counter selection with sucrose. All  $\beta$ -galactosidase expressing colonies (blue) contained the introduced BamH1 restriction site (upper panel). All white colonies displayed large

- 26 -

rearrangements and no product carried the diagnostic 1.5kb BamH1 restriction fragment (lower panel).

### Figure 11

Transference of ET cloning into a *recBC* + host to modify a large episome. a Scheme of the plasmid, pBAD-ET $\gamma$ , which carries the mobile ET system, and the strategy employed to target the Hoxa P1 episome. pBAD-ET $\gamma$  is based on pBAD24 and includes (i) the truncated *recE* gene (*t-recE*) under the arabinose-inducible *P*<sub>BAD</sub> promoter; (ii) the *recT* gene under the EM7 promoter; and (iii) the *red $\gamma$*  gene under the Tn5 promoter. It was transformed into NS3145, a *recA* E.coli strain which contained the Hoxa P1 episome. After arabinose induction, competent cells were prepared and transformed with a PCR product carrying the chloramphenicol resistance gene (*cm*) flanked by *n* and *p* homology arms. *n* and *p* were chosen to recombine with a segment of the P1 vector. b Southern blots of Pvu II digested DNAs hybridized with a probe made from the P1 vector to visualize the recombination target site (upper panel) and a probe made from the chloramphenicol resistance gene (lower panel). Lane 1, DNA prepared from cells harboring the Hoxa P1 episome before ET cloning. Lanes 2-17, DNA prepared from 16 independent chloramphenicol resistant colonies.

### Figure 12

Comparison of ET cloning using the *recE/recT* genes in pBAD-ET $\gamma$  with *red $\alpha$ /red $\beta$*  genes in pBAD- $\alpha\beta\gamma$ .

The plasmids pBAD-ET $\gamma$  or pBAD- $\alpha\beta\gamma$ , depicted, were transformed into the E.coli *recA*-, *recBC* + strain, DK1 and targeted by a chloramphenicol gene as described in Fig.6 to evaluate ET cloning efficiencies. Arabinose induction of protein expression was for 1 hour.

- 27 -

**Figure 13A**

The plasmid pBAD-ETy is shown by diagram.

**Figure 13B**

The nucleic acid sequence and the protein coding portions of pBAD-ETy are depicted.

**Figure 14A**

The plasmid pBAD- $\alpha\beta\gamma$  is shown by diagram. This plasmid substantially corresponds to the plasmid shown in Fig.13 except that the recE and recT genes are substituted by the red $\alpha$  and red $\beta$  genes.

**Figure 14B**

The nucleic acid sequence and the protein coding portions of pBAD- $\alpha\beta\gamma$  are depicted.

**1. Methods****1.1. Preparation of linear fragments**

Standard PCR reaction conditions were used to amplify linear DNA fragments. The sequences of the primers used are depicted in Table 1.

**Table 1**

The Tn5-neo gene from pJP5603 (Penfold and Pemberton, Gene 118 (1992), 145-146) was amplified by using oligo pairs a/b and c/d. The chloramphenicol (cm) resistant gene from pMAK705 (Hashimoto-Gotoh and

Sekiguchi, J. *Bacteriol.* 131 (1977), 405-412) was amplified by using primer pairs e/f and n/p. The Tn5-neo gene flanked by FRT or loxP sites was amplified from pKaZ or pKaX (<http://www.embl-heidelberg.de/ExternalInfo/stewart>) using oligo pairs i/h, g/h and j/k. The sacB-neo cassette from pIB279 (Blomfield et al., *Mol. Microbiol.* 5 (1991), 1447-1457) was amplified by using oligo pair l/m. The lacZ gene fragment from pSVpaZ11 (Buchholz et al., *Nucleic Acids Res.* 24 (1996), 4256-4262) was amplified using oligo pair l'/m'. PCR products were purified using the QIAGEN PCR Purification Kit and eluted with H<sub>2</sub>O<sub>2</sub>, followed by digestion of any residual template DNA with Dpn I. After digestion, PCR products were extracted once with Phenol:CHCl<sub>3</sub>, ethanol precipitated and resuspended in H<sub>2</sub>O at approximately 0.5 µg/µl.

## 1.2 Preparation of competent cells and electroporation

Saturated overnight cultures were diluted 50 fold into LB medium, grown to an OD<sub>600</sub> of 0.5, following by chilling on ice for 15 min. Bacterial cells were centrifuged at 7,000 rpm for 10 min at 0°C. The pellet was resuspended in ice-cold 10% glycerol and centrifuged again (7,000 rpm, -5°C, 10 min). This was repeated twice more and the cell pellet was suspended in an equal volume of ice-cold 10% glycerol. Aliquots of 50 µl were frozen in liquid nitrogen and stored at -80°C. Cells were thawed on ice and 1 µl DNA solution (containing, for co-transformation, 0.3 µg plasmid and 0.2 µg PCR products; or, for transformation, 0.2 µg PCR products) was added. Electroporation was performed using ice-cold cuvettes and a Bio-Rad Gene Pulser set to 25 µFD, 2.3 kV with Pulse Controller set at 200 ohms. LB medium (1 ml) was added after electroporation. The cells were incubated at 37°C for 1 hour with shaking and then spread on antibiotic plates.

## 1.3 Induction of RecE and RecT expression



- 29 -

E.coli JC5547 carrying pBAD24-recET was cultured overnight in LB medium plus 0.2% glucose, 100  $\mu$ g/ml ampicillin. Five parallel LB cultures, one of which (0) included 0.2% glucose, were started by a 1/100 inoculation. The cultures were incubated at 37°C with shaking for 4 hours and 0.1% L-arabinose was added 3, 2, 1 or 1/2 hour before harvesting and processing as above. Immediately before harvesting, 100  $\mu$ l was removed for analysis on a 10% SDS-polyacrylamide gel. E.coli NS3145 carrying Hoxa-P1 and pBAD-ETy was induced by 0.1% L-arabinose for 90 min before harvesting.

#### 1.4 Transient transformation of FLP and Cre expression plasmids

The FLP and Cre expression plasmids, 705-Cre and 705-FLP (Buchholz et al, Nucleic Acids Res. 24 (1996), 3118-3119), based on the pSC101 temperature sensitive origin, were transformed into rubidium chloride competent bacterial cells. Cells were spread on 25  $\mu$ g/ml chloramphenicol plates, and grown for 2 days at 30°C, whereupon colonies were picked, replated on L-agar plates without any antibiotics and incubated at 40°C overnight. Single colonies were analyzed on various antibiotic plates and all showed the expected loss of chloramphenicol and kanamycin resistance.

#### 1.5 Sucrose counter selection of sacB expression

The E.coli JC9604lacZ strain, generated as described in Fig.11, was cotransformed with a sacB-neo PCR fragment and pSVpaX1 (Buchholz et al, Nucleic Acids Res. 24 (1996), 4256-4262). After selection on 100  $\mu$ g/ml ampicillin, 50  $\mu$ g/ml kanamycin plates, pSVpaX-sacB-neo plasmids were isolated and cotransformed into fresh JC9604lacZ cells with a PCR fragment amplified from pSVpaX1 using primers l'/m'. Oligo m' carried a silent point mutation which generated a BamHI site. Cells were plated on 7% sucrose, 100  $\mu$ g/ml ampicillin, 40  $\mu$ g/ml X-gal plates and incubated at

28°C for 2 days. The blue and white colonies grown on sucrose plates were counted and further checked by restriction analysis.

## 1.6 Other methods

DNA preparation and Southern analysis were performed according to standard procedures. Hybridization probes were generated by random priming of fragments isolated from the Tn5 neo gene (PvuII), Hoxa3 gene (both HindIII fragments), lacZ genes (EcoRI and BamHI fragments from pSVpaX1), cm gene (BstBI fragments from pMAK705) and P1 vector fragments (2.2 kb EcoRI fragments from P1 vector).

## 2. Results

### 2.1 Identification of recombination events in E.coli

To identify a flexible homologous recombination reaction in E.coli, an assay based on recombination between linear and circular DNAs was designed (Fig.1, Fig.3). Linear DNA carrying the Tn5 kanamycin resistance gene (neo) was made by PCR (Fig.3a). Initially, the oligonucleotides used for PCR amplification of neo were 60mers consisting of 42 nucleotides at their 5' ends identical to chosen regions in the plasmid and, at the 3' ends, 18 nucleotides to serve as PCR primers. Linear and circular DNAs were mixed in equimolar proportions and co-transformed into a variety of E.coli hosts. Homologous recombination was only detected in sbcA E.coli hosts. More than 95% of double ampicillin/kanamycin resistant colonies (Fig.3b) contained the expected homologously recombined plasmid as determined by restriction digestion and sequencing. Only a low background of kanamycin resistance, due to genomic integration of the neo gene, was apparent (not shown).

- 31 -

The linear plus circular recombination reaction was characterized in two ways. The relationship between homology arm length and recombination efficiency was simple, with longer arms recombining more efficiently (Fig.3c). Efficiency increased within the range tested, up to 60 bp. The effect of distance between the two chosen homology sites in the recipient plasmid was examined (Fig.3d). A set of eight PCR fragments was generated by use of a constant left homology arm with differing right homology arms. The right homology arms were chosen from the plasmid sequence to be 0 - 3100 bp from the left. Correct products were readily obtained from all, with less than 4 fold difference between them, although the insertional product (0) was least efficient. Correct products also depended on the presence of both homology arms, since PCR fragments containing only one arm failed to work.

## 2.2 Involvement of RecE and RecT

The relationship between host genotype and this homologous recombination reaction was more systematically examined using a panel of E.coli strains deficient in various recombination components (Table 2).

Table 2

Only the two sbcA strains, JC8679 and JC9604 presented the intended recombination products and RecA was not required. In sbcA strains, expression of RecE and RecT is activated. Dependence on recE can be inferred from comparison of JC8679 with JC8691. Notably no recombination products were observed in JC9387 suggesting that the sbcBC background is not capable of supporting homologous recombination based on 50 nucleotide homology arms.

To demonstrate that RecE and RecT are involved, part of the recET operon was cloned into an inducible expression vector to create pBAD24-recET

(Fig.6a). the *recE* gene was truncated at its N-terminal end, as the first 588 a.a.s of RecE are dispensable. The *recBC* strain, JC5547, was transformed with pBAD24-*recET* and a time course of RecE/RecT induction performed by adding arabinose to the culture media at various times before harvesting for competent cells. The batches of harvested competent cells were evaluated for protein expression by gel electrophoresis (Fig.6b) and for recombination between a linear DNA fragment and the endogenous pBAD24-*recET* plasmid (Fig.6c). Without induction of RecE/RecT, no recombinant products were found, whereas recombination increased in approximate concordance with increased RecE/RecT expression. This experiment also shows that co-transformation of linear and circular DNAs is not essential and the circular recipient can be endogenous in the host. From the results shown in Figs.3, 6 and Table 2, we conclude that RecE and RecT mediate a very useful homologous recombination reaction in *recBC* *E.coli* at workable frequencies. Since RecE and RecT are involved, we refer to this way of recombining linear and circular DNA fragments as "ET cloning".

### 2.3 Application of ET cloning to large target DNAs

To show that large DNA episomes could be manipulated in *E.coli*, a > 76 kb P1 clone that contains at least 59 kb of the intact mouse *Hoxa* complex, (confirmed by DNA sequencing and Southern blotting), was transferred to an *E.coli* strain having an *sbvA* background (JC9604) and subjected to two rounds of ET cloning. In the first round, the Tn903 kanamycin resistance gene resident in the P1 vector was replaced by an ampicillin resistance gene (Fig.4). In the second round, the interval between the *Hoxa3* and *a4* genes was targeted either by inserting the *neo* gene between two base pairs upstream of the *Hoxa3* proximal promoter, or by deleting 6203 bp between the *Hoxa3* and *a4* genes (Fig.8a). Both insertional and deletional ET cloning products were readily obtained (Fig.8b, lanes 2, 3 and 5) showing that the

- 33 -

two rounds of ET cloning took place in this large E.coli episome with precision and no apparent unintended recombination.

The general applicability of ET cloning was further examined by targeting a gene in the E.coli chromosome (Fig.9a). The  $\beta$ -galactosidase (*lacZ*) gene of JC9604 was chosen so that the ratio between correct and incorrect recombinants could be determined by evaluating  $\beta$ -galactosidase expression. Standard conditions (0.2  $\mu$ g PCR fragment; 50  $\mu$ l competent cells), produced 24 primary colonies, 20 of which were correct as determined by  $\beta$ -galactosidase expression (Fig.9b), and DNA analysis (Fig.9c, lanes 3-6).

#### 2.4 Secondary recombination reactions to remove operational sequences

The products of ET cloning as described above are limited by the necessary inclusion of selectable marker genes. Two different ways to use a further recombination step to remove this limitation were developed. In the first way, site specific recombination mediated by either Flp or Cre recombinase was employed. In the experiments of Figs.8 and 9, either Flp recombination target sites (FRTs) or Cre recombination target sites (loxPs) were included to flank the neo gene in the linear substrates. Recombination between the FRTs or loxPs was accomplished by Flp or Cre, respectively, expressed from plasmids with the pSC101 temperature sensitive replication origin (Hashimoto-Gotoh and Sekiguchi, J.Bacteriol. 131 (1977), 405-412) to permit simple elimination of these plasmids after site specific recombination by temperature shift. The precisely recombined Hoxa P1 vector was recovered after both ET and Flp recombination with no other recombination products apparent (Fig.8, lanes 4 and 6). Similarly, Cre recombinase precisely recombined the targeted *lacZ* allele (Fig.9, lanes 7-10). Thus site specific recombination can be readily coupled with ET cloning to remove operational sequences and leave a 34 bp site specific recombination target site at the point of DNA manipulation.

In the second way to remove the selectable marker gene, two rounds of ET cloning, combining positive and counter selection steps, were used to leave the DNA product free of any operational sequences (Fig.10a).

Additionally this experiment was designed to evaluate, by a functional test based on  $\beta$ -galactosidase activity, whether ET cloning promoted small mutations such as frame shift or point mutations within the region being manipulated. In the first round, the lacZ gene of pSVpaX1 was disrupted with a 3.3 kb PCR fragment carrying the neo and B.subtilis sacB (Blomfield et al., Mol.Microbiol. 5 (1991), 1447-1457) genes, by selection for kanamycin resistance (Fig.10a). As shown above for other positively selected recombination products, virtually all selected colonies were white (Fig.10b), indicative of successful lacZ disruption, and 17 of 17 were confirmed as correct recombinants by DNA analysis. In the second round, a 1.5 kb PCR fragment designed to repair lacZ was introduced by counter selection against the sacB gene. Repair of lacZ included a silent point mutation to create a BamH1 restriction site. Approximately one quarter of sucrose resistant colonies expressed  $\beta$ -galactosidase, and all analyzed (17 of 17; Fig.10c) carried the repaired lacZ gene with the BamH1 point mutation. The remaining three quarters of sucrose resistant colonies did not express  $\beta$ -galactosidase, and all analyzed (17 of 17; Fig.10c) had undergone a variety of large mutational events, none of which resembled the ET cloning product. Thus, in two rounds of ET cloning directed at the lacZ gene, no disturbances of  $\beta$ -galactosidase activity by small mutations were observed, indicating the RecE/RecT recombination works with high fidelity. The significant presence of incorrect products observed in the counter selection step is an inherent limitation of the use of counter selection, since any mutation that ablates expression of the counter selection gene will be selected. Notably, all incorrect products were large mutations and therefore easily distinguished from the correct ET product by DNA analysis. In a different experiment (Fig.5), we observed that ET cloning into pZero2.1 (InVitroGen) by counter selection against the ccdB gene gave

- 35 -

a lower background of incorrect products (8%), indicating that the counter selection background is variable according to parameters that differ from those that influence ET cloning efficiencies.

## 2.5 Transference of ET cloning between E.coli hosts

The experiments shown above were performed in recBC- E.coli hosts since the sbcA mutation had been identified as a suppressor of recBC (Barbour et al., Proc.Natl.Acad.Sci. USA 67 (1970), 128-135; Clark, Genetics 78 (1974), 259-271). However, many useful E.coli strains are recBC+, including strains commonly used for propagation of P1, BAC or PAC episomes. To transfer ET cloning into recBC+ strains, we developed pBAD-ET $\gamma$  and pBAD- $\sigma\beta\gamma$  (Figs.13 and 14). These plasmids incorporate three features important to the mobility of ET cloning. First, RecBC is the major E.coli exonuclease and degrades introduced linear fragments. Therefore the RecBC inhibitor, Red $\gamma$  (Murphy, J.Bacteriol. 173 (1991), 5808-5821), was included. Second, the recombinogenic potential of RecE/RecT, or Red $\alpha$ /Red $\beta$ , was regulated by placing recE or red $\alpha$  under an inducible promoter. Consequently ET cloning can be induced when required and undesired recombination events which are restricted at other times. Third, we observed that ET cloning efficiencies are enhanced when RecT, or Red $\beta$ , but not RecE, or Red $\alpha$ , is overexpressed. Therefore we placed recT, or red $\beta$ , under the strong, constitutive, EM7 promoter.

pBAD-ET $\gamma$  was transformed into NS3145 E.coli harboring the original Hoxa P1 episome (Fig.11a). A region in the P1 vector backbone was targeted by PCR amplification of the chloramphenicol resistance gene (cm) flanked by n and p homology arms. As described above for positively selected ET cloning reactions, most (> 90%) chloramphenicol resistant colonies were correct. Notably, the overall efficiency of ET cloning, in terms of linear DNA transformed, was nearly three times better using pBAD-ET $\gamma$  than with similar experiments based on targeting the same episome in the sbcA host,

JC9604. This is consistent with our observation that overexpression of RecT improves ET cloning efficiencies.

5 A comparison between ET cloning efficiencies mediated by RecE/RecT, expressed from pBAD-ET $\gamma$ , and Red $\alpha$ /Red $\beta$ , expressed from pBAD- $\alpha\beta\gamma$  was made in the recA-, recBC + E.coli strain, DK1 (Fig.12). After transformation of E.coli DK1 with either pBAD-ET $\gamma$  or pBAD- $\alpha\beta\gamma$ , the same experiment as described in Figure 6a.c. to replace the bla gene of the pBAD vector with a chloramphenicol gene was performed. Both pBAD-ET $\gamma$  or pBAD- $\alpha\beta\gamma$   
10 presented similar ET cloning efficiencies in terms of responsiveness to arabinose induction of RecE and Red $\alpha$ , and number of targeted events.



Table 2

E.coli Strains	Genotypes	Amp+Kan	Amp
			$\times 10^8/\mu\text{g}$
JC8679	<i>recBC sbcA</i>	318	2.30
JC9604	<i>recA recBC sbcA</i>	114	0.30
JC8691	<i>recBC sbcA recE</i>	0	0.37
JC5547	<i>recA recBC</i>	0	0.37
JC5519	<i>recBC</i>	0	1.80
JC15329	<i>recA recBC sbcBC</i>	0	0.03
JC9387	<i>recBC sbcBC</i>	0	2.20
JC8111	<i>recBC sbcBC recF</i>	0	2.40
JC9366	<i>recA</i>	0	0.37
JC13031	<i>recJ</i>	0	0.45

## Claims

1. A method for cloning DNA molecules in cells comprising the steps of:
  - a) providing a host cell capable of performing homologous recombination,
  - b) contacting in said host cell a first DNA molecule which is capable of being replicated in said host cell with a second DNA molecule comprising at least two regions of sequence homology to regions on the first DNA molecule, under conditions which favour homologous recombination between said first and second DNA molecules and
  - c) selecting a host cell in which homologous recombination between said first and second DNA molecules has occurred.
2. The method according to claim 1 wherein the homologous recombination occurs via the recET cloning mechanism.
3. The method according to claim 2 wherein the host cell is capable of expressing recE and recT genes.
4. The method according to claim 3 wherein the recE and recT genes are selected from E.coli recE and recT genes or from  $\lambda$  red $\alpha$  and red $\beta$  genes.
5. The method according to claim 3 or 4 wherein the host cell is transformed with at least one vector capable of expressing recE and/or recT genes.
6. The method of claim 3, 4 or 5 wherein the expression of the recE and/or recT genes is under control of a regulatable promoter.

7. The method of claim 5 or 6 wherein the recT gene is overexpressed versus the recE gene.
8. The method according to any one of claims 3 to 7 wherein the recE gene is selected from a nucleic acid molecule comprising
  - (a) the nucleic acid sequence from position 1320 (ATG) to 2159 (GAC) as depicted in Fig.7B,
  - (b) the nucleic acid sequence from position 1320 (ATG) to 1998 (CGA) as depicted in Fig.13B,
  - (c) a nucleic acid encoding the same polypeptide within the degeneracy of the genetic code and/or
  - (d) a nucleic acid sequence which hybridizes under stringent conditions with the nucleic acid sequence from (a), (b) and/or (c).
9. The method according to any one of claims 3 to 8 wherein the recT gene is selected from a nucleic acid molecule comprising
  - (a) the nucleic acid sequence from position 2155 (ATG) to 2961 (GAA) as depicted in Fig.7B,
  - (b) the nucleic acid sequence from position 2086 (ATG) to 2868 (GCA) as depicted in Fig.13B,
  - (c) a nucleic acid encoding the same polypeptide within the degeneracy of the genetic code and/or
  - (d) a nucleic acid sequence which hybridizes under stringent conditions with the nucleic acid sequences from (a), (b) and/or (c).
10. The method according to any one of the previous claims wherein the host cell is a gram-negative bacterial cell.
11. The method according to claim 10 wherein the host cell is an Escherichia coli cell.

12. The method according to claim 11 wherein the host cell is an Escherichia coli K12 strain.
13. The method according to claim 12 wherein the E.coli strain is selected from JC 8679 and JC 9604.
14. The method according to any one of the previous claims wherein the host cell further is capable of expressing a recBC inhibitor gene.
15. The method according to claim 14 wherein the host cell is transformed with a vector expressing the recBC inhibitor gene.
16. The method according to claim 14 or 15 wherein the recBC inhibitor gene is selected from a nucleic acid molecule comprising
- (a) the nucleic acid sequence from position 3588 (ATG) to 4002 (GTA) as depicted in Fig.13B,
  - (b) a nucleic acid encoding the same polypeptide within the degeneracy of the genetic code and/or
  - (c) a nucleic acid sequence which hybridizes under stringent conditions (as defined above) with the nucleic acid sequence from (a) and/ or (b).
17. The method according to any one of claims 13 to 16 wherein the host cell is a prokaryotic recBC<sup>+</sup> cell.
18. The method according to any one of the previous claims wherein the first DNA molecule is circular.
19. The method according to any one of the previous claims wherein the first DNA molecule is an extrachromosomal DNA molecule containing an origin of replication which is operative in the host cell.

20. The method according to claim 18 or 19 wherein the first DNA molecule is selected from plasmids, cosmids, P1 vectors, BAC vectors and PAC vectors.
- 5 21. The method according to any one of claims 1-18 wherein the first DNA molecule is a host cell chromosome.
22. The method according to any one of the previous claims wherein the second DNA molecule is linear.
- 10 23. The method according to any one of the previous claims wherein the regions of sequence homology are at least 15 nucleotides each.
24. The method according to one of claims 1 to 16 wherein the second DNA molecule is obtained by an amplification reaction.
- 15 25. The method according to one of the previous claims wherein the first and/or second DNA molecules are introduced into the host cells by transformation.
- 20 26. The method according to claim 25 wherein the transformation method is electroporation.
27. The method according to one of claims 1 to 26 wherein the first and second DNA molecules are introduced into the host cell simultaneously by co-transformation.
- 25 28. The method according to one of claims 1 to 26 wherein the second DNA molecule is introduced into a host cell in which the first DNA molecule is already present.
- 30

43. A reagent kit for cloning comprising  
(a) a host cell  
(b) means of expressing *recE* and *recT* genes in said host cell and  
(c) a recipient cloning vehicle capable of being replicated in said cell.
- 5 44. The reagent kit according to claim 43 wherein the means (b) comprise a vector system capable of expressing the *recE* and *recT* genes in the host cell.
- 10 45. The reagent kit according to claim 43 or 44 wherein the *recE* and *recT* genes are selected from *E.coli recE* and *recT* genes or from  $\lambda$  *red $\alpha$*  and *red $\beta$*  genes.
- 15 46. A reagent kit for cloning comprising  
(a) a source for *RecE* and *RecT* proteins and  
(b) a recipient cloning vehicle capable of being propagated in a host cell.
- 20 47. The reagent kit according to claim 46 further comprising a host cell suitable for propagating said recipient cloning vehicle.
- 25 48. The reagent kit according to claim 46 or 47 wherein said *RecE* and *RecT* or proteins are selected from *E.coli RecE* and *RecT* proteins or from phage  $\lambda$  *Red $\alpha$*  and *Red $\beta$*  proteins.
49. The reagent kit according to any one of claims 43-48 further comprising means for expressing a site specific recombinase in said host cell.
- 30 50. The reagent kit according to any one of claims 43-49 further comprising nucleic acid amplification primers comprising a region of homology to said recipient cloning vehicle.

1/65




Figure 1

2/65

## Three ways to select recombinants






Figure 2




Figure 3

*a**b*

4/65

c



d




Figure 3

Figure 4a







Figure 4b




- Lane 1: 1 of P1-Hox clone in NS3145 original bacterial strain (Kan resistance)
- Lane 2-3: 2 of P1-Hox clones in JC9604 before homologous recombination (Kan resistance)
- Lane 4-6: 3 of P1-Hox clones in JC9604 after homologous recombination (Amp resistance)

Figure 5

*a**b*


8/65

Figure 6



SUBSTITUTE SHEET (RULE 26)

Figure 6



10/65

C





Figure 6



Figure 7a



t-recE --- truncated recE (from 588 aa ---> end. 866 aa)

12/65

Figure 7b

```

1  ATCGATGCATAATGTGCCTGTCAAATGGACGAAGCAGGGATTCT
44  TGCAAACCCTATGCTACTCCGTCAAGCCGTCAATTGTCTGATT
87  CGTTACCAA TTA TGA CAA CTT GAC GGC TAC ATC
    293◀••• Ser Leu Lys Val Ala Val Asp

120  ATT CAC TTT TTC TTC ACA ACC GGC ACG GAA CTC
285◀Asn Val Lys Glu Glu Cys Gly Ala Arg Phe Glu

153  GCT CGG GCT GGC CCC GGT GCA TTT TTT AAA TAC
274◀Ser Pro Ser Ala Gly Thr Cys Lys Lys Phe Val

186  CCG CGA GAA ATA GAG TTG ATC GTC AAA ACC AAC
263◀Arg Ser Phe Tyr Leu Gln Asp Asp Phe Gly Val

219  ATT GCG ACC GAC GGT GGC GAT AGG CAT CCG GGT
252◀Asn Arg Gly Val Thr Ala Ile Pro Met Arg Thr

252  GGT GCT CAA AAG CAG CTT CGC CTG GCT GAT ACG
241◀Thr Ser Leu Leu Leu Lys Ala Gln Ser Ile Arg

285  TTG GTC CTC GCG CCA GCT TAA GAC GCT AAT CCC
230◀Gln Asp Glu Arg Trp Ser Leu Val Ser Ile Gly

318  TAA CTG CTG GCG GAA AAG ATG TGA CAG ACG CGA
219◀Leu Gln Gln Arg Phe Leu His Ser Leu Arg Ser

351  CGG CGA CAA GCA AAC ATG CTG TGC GAC GCT GGC
208◀Pro Ser Leu Cys Val His Gln Ala Val Ser Ala
    EcoRV

384  GAT ATC AAA ATT GCT GTC TGC CAG GTG ATC GCT
197◀Ile Asp Phe Asn Ser Asp Ala Leu His Asp Ser

417  GAT GTA CTG ACA AGC CTC GCG TAC CCG ATT ATC
186◀Ile Tyr Gln Cys Ala Glu Arg Val Arg Asn Asp

```

SUBSTITUTE SHEET (RULE 26)

13/65

Figure 7b (cont'd)

450	CAT	CGG	TGG	ATG	GAG	CGA	CTC	GTT	AAT	CGC	TTC
175	Met	Pro	Pro	His	Leu	Ser	Glu	Asn	Ile	Ala	Glu
483	CAT	GCG	CCG	CAG	TAA	CAA	TTG	CTC	AAG	CAG	ATT
164	Met	Arg	Arg	Leu	Leu	Leu	Gln	Glu	Leu	Leu	Asn
516	TAT	CGC	CAG	CAG	CTC	CGA	ATA	GCG	CCC	TTC	CCC
153	Ile	Ala	Leu	Leu	Glu	Ser	Tyr	Arg	Gly	Glu	Gly
549	TTG	CCC	GGC	GTT	AAT	GAT	TTG	CCC	AAA	CAG	GTC
142	Gln	Gly	Ala	Asn	Ile	Ile	Gln	Gly	Phe	Leu	Asp
582	GCT	GAA	ATG	CGG	CTG	GTG	CGC	TTC	ATC	CGG	GCG
131	Ser	Phe	His	Pro	Gln	His	Ala	Glu	Asp	Pro	Arg
615	AAA	GAA	CCC	CGT	ATT	GGC	AAA	TAT	TGA	CGG	CCA
120	Phe	Phe	Gly	Thr	Asn	Ala	Phe	Ile	Ser	Pro	Trp
648	GTT	AAG	CCA	TTC	ATG	CCA	GTA	GGC	GCG	CGG	ACG
109	Asn	Leu	Trp	Glu	His	Trp	Tyr	Ala	Arg	Pro	Arg
681	AAA	GTA	AAC	CCA	CTG	GTG	ATA	CCA	TTC	GCG	AGC
98	Phe	Tyr	Val	Trp	Gln	His	Tyr	Trp	Glu	Arg	Ala
714	CTC	CGG	ATG	ACG	ACC	GTA	GTG	ATG	AAT	CTC	TCC
87	Glu	Pro	His	Arg	Gly	Tyr	His	His	Ile	Glu	Gly
747	TGG	CGG	GAA	CAG	CAA	AAT	ATC	ACC	CGG	TCG	GCA
76	Pro	Pro	Phe	Leu	Leu	Ile	Asp	Gly	Pro	Arg	Cys
780	AAC	AAA	TTC	TCG	TCC	CTG	ATT	TTT	CAC	CAC	CCC
65	Val	Phe	Glu	Arg	Gly	Gln	Asn	Lys	Val	Val	Gly
813	CTG	ACC	GCG	AAT	GGT	GAG	ATT	GAG	AAT	ATA	ACC
54	Gln	Gly	Arg	Ile	Thr	Leu	Asn	Leu	Ile	Tyr	Gly
846	TTT	CAT	TCC	CAG	CGG	TCG	GTC	GAT	AAA	AAA	ATC
43	Lys	Met	Gly	Leu	Pro	Arg	Asp	Ile	Phe	Phe	Asp

14/65

Figure 7b (cont'd)

879 GAG ATA ACC GTT GGC CTC AAT CGG CGT TAA ACC  
 32◀ Leu Tyr Gly Asn Ala Glu Ile Pro Thr Leu Gly  
 912 CGC CAC CAG ATG GGC ATT AAA CGA GTA TCC CGG  
 21◀ Ala Val Leu His Ala Asn Phe Ser Tyr Gly Pro  
 945 CAG CAG GGG ATC ATT TTG CGC TTC AGC CAT  
 10◀ Leu Leu Pro Asp Asn Gln Ala Glu Ala Met  
 975 ACTTTTCATA CTCCCGCCAT TCAGAGAAGA AACCAATTGT  
 1015 CCATATTGCA TCAGACATTG CCGTCACTGC GTCTTTTACT  
 1055 GGCTCTTCTC GCTAACC AAA CCGGTAACCC ~~CG~~CTTATTAA  
 1095 AAGCATTTCTG TAACAAAGCG GGACCAAAGC CATGACAAAA  
 1135 ACGCGTAACA AAAGTGTCTA TAATCACGGC AGAAAAGTCC  
 1175 ACATTGATTA TTTGCACGGC GTCACACTTT GCTATGCCAT  
 BamHI  
 1215 AGCATTTTTA TCCATAAGAT TAGCGGATCC TACCTGACGC  
 1255 TTTTATATCGC AACTCTCTAC TGTTTCTCCA TACCCGTTTT  
 NheI EcoRI NcoI  
 1295 TTTGGGCTAG CAGGAGGAAT TCACC ATG GAT CCC GTA  
 1▶ Met Asp Pro Val  
 1332 ATC GTA GAA GAC ATA GAG CCA GGT ATT TAT TAC  
 5▶ Ile Val Glu Asp Ile Glu Pro Gly Ile Tyr Tyr  
 1365 GGA ATT TCG AAT GAG AAT TAC CAC GCG GGT CCC  
 16▶ Gly Ile Ser Asn Glu Asn Tyr His Ala Gly Pro  
 1398 GGT ATC AGT AAG TCT CAG CTC GAT GAC ATT GCT

15/65

Figure 7b (cont'd)

27▶	Gly	Ile	Ser	Lys	Ser	Gln	Leu	Asp	Asp	Ile	Ala
1431	GAT	ACT	CCG	GCA	CTA	TAT	TTG	TGG	CGT	AAA	AAT
38▶	Asp	Thr	Pro	Ala	Leu	Tyr	Leu	Trp	Arg	Lys	Asn
1464	GCC	CCC	GTG	GAC	ACC	ACA	AAG	ACA	AAA	ACG	CTC
49▶	Ala	Pro	Val	Asp	Thr	Thr	Lys	Thr	Lys	Thr	Leu
1497	GAT	TTA	GGA	ACT	GCT	TTC	CAC	TGC	CGG	GTA	CTT
60▶	Asp	Leu	Gly	Thr	Ala	Phe	His	Cys	Arg	Val	Leu
				EcoRI							
1530	GAA	CCG	GAA	GAA	TTC	AGT	AAC	CGC	TTT	ATC	GTA
71▶	Glu	Pro	Glu	Glu	Phe	Ser	Asn	Arg	Phe	Ile	Val
1563	GCA	CCT	GAA	TTT	AAC	CGC	CGT	ACA	AAC	GCC	GGA
82▶	Ala	Pro	Glu	Phe	Asn	Arg	Arg	Thr	Asn	Ala	Gly
1596	AAA	GAA	GAA	GAG	AAA	GCG	TTT	CTG	ATG	GAA	TGC
93▶	Lys	Glu	Glu	Glu	Lys	Ala	Phe	Leu	Met	Glu	Cys
1629	GCA	AGC	ACA	GGA	AAA	ACG	GTT	ATC	ACT	GCG	GAA
104▶	Ala	Ser	Thr	Gly	Lys	Thr	Val	Ile	Thr	Ala	Glu
1662	GAA	GGC	CGG	AAA	ATT	GAA	CTC	ATG	TAT	CAA	AGC
115▶	Glu	Gly	Arg	Lys	Ile	Glu	Leu	Met	Tyr	Gln	Ser

16/65

Figure 7b (cont'd)

1695	GTT	ATG	GCT	TTG	CCG	CTG	GGG	CAA	TGG	CTT	GTT
126▶	Val	Met	Ala	Leu	Pro	Leu	Gly	Gln	Trp	Leu	Val
1728	GAA	AGC	GCC	GGA	CAC	GCT	GAA	TCA	TCA	ATT	TAC
137▶	Glu	Ser	Ala	Gly	His	Ala	Glu	Ser	Ser	Ile	Tyr
1761	TGG	GAA	GAT	CCT	GAA	ACA	GGA	ATT	TTG	TGT	CGG
148▶	Trp	Glu	Asp	Pro	Glu	Thr	Gly	Ile	Leu	Cys	Arg
1794	TGC	CGT	CCG	GAC	AAA	ATT	ATC	CCT	GAA	TTT	CAC
159▶	Cys	Arg	Pro	Asp	Lys	Ile	Ile	Pro	Glu	Phe	His
1827	TGG	ATC	ATG	GAC	GTG	AAA	ACT	ACG	GCG	GAT	ATT
170▶	Trp	Ile	Met	Asp	Val	Lys	Thr	Thr	Ala	Asp	Ile
1860	CAA	CGA	TTC	AAA	ACC	GCT	TAT	TAC	GAC	TAC	CGC
181▶	Gln	Arg	Phe	Lys	Thr	Ala	Tyr	Tyr	Asp	Tyr	Arg
1893	TAT	CAC	GTT	CAG	GAT	GCA	TTC	TAC	AGT	GAC	GGT
192▶	Tyr	His	Val	Gln	Asp	Ala	Phe	Tyr	Ser	Asp	Gly
1926	TAT	GAA	GCA	CAG	TTT	GGA	GTG	CAG	CCA	ACT	TTC
203▶	Tyr	Glu	Ala	Gln	Phe	Gly	Val	Gln	Pro	Thr	Phe
1959	GTT	TTT	CTG	GTT	GCC	AGC	ACA	ACT	ATT	GAA	TGC
214▶	Val	Phe	Leu	Val	Ala	Ser	Thr	Thr	Ile	Glu	Cys
1992	GGA	CGT	TAT	CCG	GTT	GAA	ATT	TTC	ATG	ATG	GGC

SUBSTITUTE SHEET (RULE 26)

17/65

Figure 7b (cont'd)

225▶ Gly Arg Tyr Pro Val Glu Ile Phe Met Met Gly  
 2025 GAA GAA GCA AAA CTG GCA GGT CAA CAG GAA TAT

236▶ Glu Glu Ala Lys Leu Ala Gly Gln Gln Glu Tyr  
 2058 CAC CGC AAT CTG CGA ACC CTG TCT GAC TGC CTG

247▶ His Arg Asn Leu Arg Thr Leu Ser Asp Cys Leu  
 Ball  
 2091 AAT ACC GAT GAA TGG CCA GCT ATT AAG ACA TTA

258▶ Asn Thr Asp Glu Trp Pro Ala Ile Lys Thr Leu  
 2124 TCA CTG CCC CGC TGG GCT AAG GAA TAT GCAA

269▶ Ser Leu Pro Arg Trp Ala Lys Glu Tyr AlaA  
 2155 ATG ACT AAG CAA CCA CCA ATC GCA AAA GCC GAT  
 1▶ Met Thr Lys Gln Pro Pro Ile Ala Lys Ala Asp  
 279▶ s nAs p . . .

2188 CTG CAA AAA ACT CAG GGA AAC CGT GCA CCA GCA  
 12▶ Leu Gln Lys Thr Gln Gly Asn Arg Ala Pro Ala  
 2221 GCA GTT AAA AAT AGC GAC GTG ATT AGT TTT ATT  
 23▶ Ala Val Lys Asn Ser Asp Val Ile Ser Phe Ile  
 2254 AAC CAG CCA TCA ATG AAA GAG CAA CTG GCA GCA  
 34▶ Asn Gln Pro Ser Met Lys Glu Gln Leu Ala Ala  
 Ndel  
 2287 GCT CTT CCA CGC CAT ATG ACG GCT GAA CGT ATG  
 45▶ Ala Leu Pro Arg His Met Thr Ala Glu Arg Met

18/65

Figure 7b (cont'd)

2320	ATC	CGT	ATC	GCC	ACC	ACA	GAA	ATT	CGT	AAA	GTT
56▶	Ile	Arg	Ile	Ala	Thr	Thr	Glu	Ile	Arg	Lys	Val
2353	CCG	GCG	TTA	GGA	AAC	TGT	GAC	ACT	ATG	AGT	TTT
67▶	Pro	Ala	Leu	Gly	Asn	Cys	Asp	Thr	Met	Ser	Phe
2386	GTC	AGT	GCG	ATC	GTA	CAG	TGT	TCA	CAG	CTC	GGA
78▶	Val	Ser	Ala	Ile	Val	Gln	Cys	Ser	Gln	Leu	Gly
2419	CTT	GAG	CCA	GGT	AGC	GCC	CTC	GGT	CAT	GCA	TAT
89▶	Leu	Glu	Pro	Gly	Ser	Ala	Leu	Gly	His	Ala	Tyr
2452	TTA	CTG	CCT	TTT	GGT	AAT	AAA	AAC	GAA	AAG	AGC
100▶	Leu	Leu	Pro	Phe	Gly	Asn	Lys	Asn	Glu	Lys	Ser
2485	GGT	AAA	AAG	AAC	GTT	CAG	CTA	ATC	ATT	GGC	TAT
111▶	Gly	Lys	Lys	Asn	Val	Gln	Leu	Ile	Ile	Gly	Tyr
2518	CGC	GGC	ATG	ATT	GAT	CTG	GCT	CGC	CGT	TCT	GGT
122▶	Arg	Gly	Met	Ile	Asp	Leu	Ala	Arg	Arg	Ser	Gly
2551	CAA	ATC	GCC	AGC	CTG	TCA	GCC	CGT	GTT	GTC	CGT
133▶	Gln	Ile	Ala	Ser	Leu	Ser	Ala	Arg	Val	Val	Arg
2584	GAA	GGT	GAC	GAG	TTT	AGC	TTC	GAA	TTT	GGC	CTT
144▶	Glu	Gly	Asp	Glu	Phe	Ser	Phe	Glu	Phe	Gly	Leu
2617	GAT	GAA	AAG	TTA	ATA	CAC	CGC	CCG	GGA	GAA	AAC
155▶	Asp	Glu	Lys	Leu	Ile	His	Arg	Pro	Gly	Glu	Asn
2650	GAA	GAT	GCC	CCG	GTT	ACC	CAC	GTC	TAT	GCT	GTC
166▶	Glu	Asp	Ala	Pro	Val	Thr	His	Val	Tyr	Ala	Val
2683	GCA	AGA	CTG	AAA	GAC	GGA	GGT	ACT	CAG	TTT	GAA
177▶	Ala	Arg	Leu	Lys	Asp	Gly	Gly	Thr	Gln	Phe	Glu
2716	GTT	ATG	ACG	CGC	AAA	CAG	ATT	GAG	CTG	GTG	CGC
188▶	Val	Met	Thr	Arg	Lys	Gln	Ile	Glu	Leu	Val	Arg

SUBSTITUTE SHEET (RULE 26)



19/65

Figure 7b (cont'd)

2749 AGC CTG AGT AAA GCT GGT AAT AAC GGG CCG TGG  
 199► Ser Leu Ser Lys Ala Gly Asn Asn Gly Pro Trp  
 2782 GTA ACT CAC TGG GAA GAA ATG GCA AAG AAA ACG  
 210► Val Thr His Trp Glu Glu Met Ala Lys Lys Thr  
 2815 GCT ATT CGT CGC CTG TTC AAA TAT TTG CCC GTA  
 221► Ala Ile Arg Arg Leu Phe Lys Tyr Leu Pro Val  
 2848 TCA ATT GAG ATC CAG CGT GCA GTA TCA ATG GAT  
 232► Ser Ile Glu Ile Gln Arg Ala Val Ser Met Asp  
 PstI  
 2881 GAA AAG GAA CCA CTG ACA ATC GAT CCT GCA GAT  
 243► Glu Lys Glu Pro Leu Thr Ile Asp Pro Ala Asp  
 2914 TCC TCT GTA TTA ACC GGG GAA TAC AGT GTA ATC  
 254► Ser Ser Val Leu Thr Gly Glu Tyr Ser Val Ile  
 BglII HindIII  
 2947 GAT AAT TCA GAG GAA TAG ATCTAAGCTT  
 265► Asp Asn Ser Glu Glu ...  
 2975 GGCTGTTTTG GCGGATGAGA GAAGATTTTC AGCCTGATAC  
 3015 AGATTAAATC AGAACGCAGA AGCGGTCTGA TAAACAGAA  
 3055 TTTGCCTGGC GGCAGTAGCG CGGTGGTCCC ACCTGACCCC  
 3095 ATGCCGA ACT CAGAAGTGAA ACGCCGTAGC GCCGATGGTA  
 3135 GTGTGGGGTC TCCCCATGCG AGAGTAGGGA ACTGCCAGGC  
 3175 ATCAAATAAA ACGAAAGGCT CAGTCGAAAG ACTGGGCCTT  
 3215 TCGTTTTATC TGTTGTTTGT CGGTGAACGC TCTCCTGAGT  
 3255 AGGACAAATC CGCCGGGAGC GGATTTGAAC GTTGCGAAGC  
 3295 AACGGCCCGG AGGGTGGCGG GCAGGACGCC CGCCATAAAC  
 3335 TGCCAGGCAT CAAATTAAGC AGAAGGCCAT CCTGACGGAT

20/65

Figure 7b (cont'd)

3375 GGCCTTTTGTG CGTTTCTACA AACTCTTTTG TTTATTTTTC  
 3415 TAAATACATT CAAATATGTA TCCGCTCATG AGACAATAAC  
 3455 CCTGATAAAT GCTTCAATAA TATTGAAAAA GGAAGAGT AT  
 1► Me  
 3495 G AGT ATT CAA CAT TTC CGT GTC GCC CTT ATT  
 1► t Ser Ile Gln His Phe Arg Val Ala Leu Ile  
 3526 CCC TTT TTT GCG GCA TTT TGC CTT CCT GTT TTT  
 12► Pro Phe Phe Ala Ala Phe Cys Leu Pro Val Phe  
 3559 GCT CAC CCA GAA ACG CTG GTG AAA GTA AAA GAT  
 23► Ala His Pro Glu Thr Leu Val Lys Val Lys Asp  
 3592 GCT GAA GAT CAG TTG GGT GCA CGA GTG GGT TAC  
 34► Ala Glu Asp Gln Leu Gly Ala Arg Val Gly Tyr  
 3625 ATC GAA CTG GAT CTC AAC AGC GGT AAG ATC CTT  
 45► Ile Glu Leu Asp Leu Asn Ser Gly Lys Ile Leu  
 3658 GAG AGT TTT CGC CCC GAA GAA CGT TTT CCA ATG  
 56► Glu Ser Phe Arg Pro Glu Glu Arg Phe Pro Met  
 3691 ATG AGC ACT TTT AAA GTT CTG CTA TGT GGC GCG  
 67► Met Ser Thr Phe Lys Val Leu Leu Cys Gly Ala  
 3724 GTA TTA TCC CGT GTT GAC GCC GGG CAA GAG CAA  
 78► Val Leu Ser Arg Val Asp Ala Gly Gln Glu Gln  
 3757 CTC GGT CGC CGC ATA CAC TAT TCT CAG AAT GAC  
 89► Leu Gly Arg Arg Ile His Tyr Ser Gln Asn Asp  
 Scal  
 3790 TTG GTT GAG TAC TCA CCA GTC ACA GAA AAG CAT  
 100► Leu Val Glu Tyr Ser Pro Val Thr Glu Lys His  
 3823 CTT ACG GAT GGC ATG ACA GTA AGA GAA TTA TGC  
 111► Leu Thr Asp Gly Met Thr Val Arg Glu Leu Cys

SUBSTITUTE SHEET (RULE 26)

21/65

Figure 7b (cont'd)

3856	AGT	GCT	GCC	ATA	ACC	ATG	AGT	GAT	AAC	ACT	GCG
122▶	Ser	Ala	Ala	Ile	Thr	Met	Ser	Asp	Asn	Thr	Ala
3889	GCC	AAC	TTA	CTT	CTG	ACA	ACG	ATC	GGA	GGA	CCG
133▶	Ala	Asn	Leu	Leu	Leu	Thr	Thr	Ile	Gly	Gly	Pro
3922	AAG	GAG	CTA	ACC	GCT	TTT	TTG	CAC	AAC	ATG	GGG
144▶	Lys	Glu	Leu	Thr	Ala	Phe	Leu	His	Asn	Met	Gly
3955	GAT	CAT	GTA	ACT	CGC	CTT	GAT	CGT	TGG	GAA	CCG
155▶	Asp	His	Val	Thr	Arg	Leu	Asp	Arg	Trp	Glu	Pro
3988	GAG	CTG	AAT	GAA	GCC	ATA	CCA	AAC	GAC	GAG	CGT
166▶	Glu	Leu	Asn	Glu	Ala	Ile	Pro	Asn	Asp	Glu	Arg
4021	GAC	ACC	ACG	ATG	CCT	GTA	GCA	ATG	GCA	ACA	ACG
177▶	Asp	Thr	Thr	Met	Pro	Val	Ala	Met	Ala	Thr	Thr
4054	TTG	CGC	AAA	CTA	TTA	ACT	GGC	GAA	CTA	CTT	ACT
188▶	Leu	Arg	Lys	Leu	Leu	Thr	Gly	Glu	Leu	Leu	Thr
4087	CTA	GCT	TCC	CGG	CAA	CAA	TTA	ATA	GAC	TGG	ATG
199▶	Leu	Ala	Ser	Arg	Gln	Gln	Leu	Ile	Asp	Trp	Met
4120	GAG	GCG	GAT	AAA	GTT	GCA	GGA	CCA	CTT	CTG	CGC
210▶	Glu	Ala	Asp	Lys	Val	Ala	Gly	Pro	Leu	Leu	Arg
4153	TCG	GCC	CTT	CCG	GCT	GGC	TGG	TTT	ATT	GCT	GAT
221▶	Ser	Ala	Leu	Pro	Ala	Gly	Trp	Phe	Ile	Ala	Asp
4186	AAA	TCT	GGA	GCC	GGT	GAG	CGT	GGG	TCT	CGC	GGT
232▶	Lys	Ser	Gly	Ala	Gly	Glu	Arg	Gly	Ser	Arg	Gly
4219	ATC	ATT	GCA	GCA	CTG	GGG	CCA	GAT	GGT	AAG	CCC
243▶	Ile	Ile	Ala	Ala	Leu	Gly	Pro	Asp	Gly	Lys	Pro
4252	TCC	CGT	ATC	GTA	GTT	ATC	TAC	ACG	ACG	GGG	AGT
254▶	Ser	Arg	Ile	Val	Val	Ile	Tyr	Thr	Thr	Gly	Ser

22/65

Figure 7b (cont'd)

4285 CAG GCA ACT ATG GAT GAA CGA AAT AGA CAG ATC  
265► Gln Ala Thr Met Asp Glu Arg Asn Arg Gln Ile  
4318 GCT GAG ATA GGT GCC TCA CTG ATT AAG CAT TGG  
276► Ala Glu Ile Gly Ala Ser Leu Ile Lys His Trp  
4351 TAA CTGTCAGACC AAGTTTACTC ATATATACTT  
287► . . .

4384 TAGATTGATT TACGCGCCCT GTAGCGGCGC ATTAAGCGCG  
4424 GCGGGTGTGG TGGTTACGCG CAGCGTGACC GCTACACTTG  
4464 CCAGCGCCCT AGCGCCCGCT CCTTTCGCTT TCTTCCCTTC  
4504 CTTTCTCGCC ACGTTCGCCG GCTTTCCTCCG TCAAGCTCTA  
4544 AATCGGGGGC TCCCTTTAGG GTTCCGATTT AGTGCTTTAC  
4584 GGCACCTCGA CCCCÀAAAAA CTTGATTTGG GTGATGGTTC  
4624 ACGTAGTGGG CCATCGCCCT GATAGACGGT TTTTCGCCCT  
4664 TTGACGTTGG AGTCCACGTT CTTTAATAGT GGACTCTTGT  
4704 TCCAAACTTG AACAACACTC AACCCCTATCT CGGGCTATTC  
4744 TTTTGATTTA TAAGGGATTT TGCCGATTTT GGCCTATTGG  
4784 TTAAAAAATG AGCTGATTTA ACAAAAATTT AACGCGAATT  
4824 TTAACAAAAT ATTAACGTTT ACAATTTAAA AGGATCTAGG  
4864 TGAAGATCCT TTTTGATAAT CTCATGACCA AAATCCCTTA  
4904 ACGTGAGTTT TCGTTCCACT GAGCGTCAGA CCCCCTAGAA  
4944 AAGATCAAAG GATCTTCTTG AGATCCTTTT TTTCTGCGCG  
4984 TAATCTGCTG CTTGCAAACA AAAAAACCAC CGCTACCAGC  
5024 GGTGGTTTGT TTGCCGGATC AAGAGCTACC AACTCTTTTT

SUBSTITUTE SHEET (RULE 26)

23/65

Figure 7b (cont'd)


5064	CCGAAGGTAA	CTGGCTTCAG	CAGAGCGCAG	ATACCAAATA
5104	CTGTCCTTCT	AGTG TAGCCG	TAGTTAGGCC	ACCACTTCAA
5144	GAAC TCTGTA	GCACCGCCTA	CATACCTCGC	TCTGCTAATC
5184	CTGTTACCAG	TGGCTGCTGC	CAGTGGCGAT	AAGTCGTGTC
5224	TTACCGGGTT	GGACTCAAGA	CGATAGTTAC	CGGATAAGGC
5264	GCAGCGGTTCG	GGCTGAACGG	GGGGTTCGTG	CACACAGCCC
5304	AGCTTGGAGC	GAACGACCTA	CACCGAACTG	AGATACCTAC
5344	AGCGTGAGCT	ATGAGAAAGC	GCCACGCTTC	CCGAAGGGAG
5384	AAAGGCGGAC	AGGTATCCGG	TAAGCGGCAG	GGTCGGAACA
5424	GGAGAGCGCA	CGAGGGGAGCT	TCCAGGGGGA	AACGCCTGGT
5464	ATCTTTATAG	TCCTGTCTGGG	TTTCGCCACC	TCTGACTTGA
5504	GCGTCGATTT	TTGTGATGCT	CGTCAGGGGG	GCGGAGCCTA
5544	TGGAAAAACG	CCAGCAACGC	GGCCTTTTTA	CGGTTCCCTGG
5584	CCTTTTGCTG	GCCTTTTGCT	CACATGTTCT	TTCCTGCGTT
5624	ATCCCCTGAT	TCTGTGGATA	ACCGTATTAC	CGCCTTTGAG
5664	TGAGCTGATA	CCGCTCGCCG	CAGCCGAACG	ACCGAGCGCA
5704	GCGAGTCAGT	GAGCGAGGAA	GCGGAAGAGC	GCCTGATGCG
5744	GTATTTTCTC	CTTACGCATC	TGTGCGGTAT	TTCACACCGC
5784	ATAGGGTCAT	GGCTGCGCCC	CGACACCCGC	CAACACCCGC
5824	TGACGCGCCC	TGACGGGCTT	GTCTGCTCCC	GGCATCCGCT
5864	TACAGACAAG	CTGTGACCGT	CTCCGGGAGC	TGCATGTGTC
5904	AGAGGTTTTTC	ACCGTCATCA	CCGAAACGCG	CGAGGCAGCA
5944	AGGAGATGGC	GCCCAACAGT	CCCCCGGCCA	CGGGGCCTGC

24/65

Figure 7b (cont'd)


5984 CACCATACCC ACGCCGAAAC AAGCGCTCAT GAGCCCGAAG  
6024 TGGCGAGCCC GATCTTCCCC ATCGGTGATG TCGGCGATAT  
6064 AGGCGCCAGC AACCGCACCT GTGGCGCCGG TGATGCCGGC  
6104 CACGATGCGT CCGGCGTAGA GGATCTGCTC ATGTTTGACA  
6144 GCTTATC

SUBSTITUTE SHEET (RULE 26)



+

Deletion



Insertion




Figure 8 a





Figure 8 a (continuing)




27/65

Figure 8 b




28/65

Figure 9a



SUBSTITUTE SHEET (RULE 26)

Figure 9

*b**c*

1 2 3 4 5 6 7 8 9 10



Figure 10a




Figure 10

*b*

pSVpaX1


pSV-sacB-neo

pSVpaX1\*



*c*


blue clones



white clones



Figure 11a



SUBSTITUTE SHEET (RULE 26)

Figure 11 b







Figure 12



SUBSTITUTE SHEET (RULE 26)



Figure 13 a



36/65

Figure 13b

```

1  ATCGATGCATAATGTGCCTGTCAAATGGACGAAGCAGGG
40  ATTCTGCAAACCCTATGCTACTCCGTCAAGCCGTCAATT
79  GTCTGATTTCGTTACCAA TTA TGA CAA CTT GAC
    293◀••• Ser Leu Lys Val

111  GGC TAC ATC ATT CAC TTT TTC TTC ACA ACC
288◀Ala Val Asp Asn Val Lys Glu Glu Cys Gly

141  GGC ACG GAA CTC GCT CGG GCT GGC CCC GGT
278◀Ala Arg Phe Glu Ser Pro Ser Ala Gly Thr

171  GCA TTT TTT AAA TAC CCG CGA GAA ATA GAG
268◀Cys Lys Lys Phe Val Arg Ser Phe Tyr Leu

201  TTG ATC GTC AAA ACC AAC ATT GCG ACC GAC
258◀Gln Asp Asp Phe Gly Val Asn Arg Gly Val

231  GGT GGC GAT AGG CAT CCG GGT GGT GCT CAA
248◀Thr Ala Ile Pro Met Arg Thr Thr Ser Leu

261  AAG CAG CTT CGC CTG GCT GAT ACG TTG GTC
238◀Leu Leu Lys Ala Gln Ser Ile Arg Gln Asp

291  CTC GCG CCA GCT TAA GAC GCT AAT CCC TAA
228◀Glu Arg Trp Ser Leu Val Ser Ile Gly Leu

321  CTG CTG GCG GAA AAG ATG TGA CAG ACG CGA
218◀Gln Gln Arg Phe Leu His Ser Leu Arg Ser

351  CGG CGA CAA GCA AAC ATG CTG TGC GAC GCT
208◀Pro Ser Leu Cys Val His Gln Ala Val Ser

381  GGC GAT ATC AAA ATT GCT GTC TGC CAG GTG
198◀Ala Ile Asp Phe Asn Ser Asp Ala Leu His

411  ATC GCT GAT GTA CTG ACA AGC CTC GCG TAC

```

37/65

Figure 13b (cont'd)

188 ◀ Asp Ser Ile Tyr Gln Cys Ala Glu Arg Val  
 441 CCG ATT ATC CAT CGG TGG ATG GAG CGA CTC  
 178 ◀ Arg Asn Asp Met Pro Pro His Leu Ser Glu  
 471 GTT AAT CGC TTC CAT GCG CCG CAG TAA CAA  
 168 ◀ Asn Ile Ala Glu Met Arg Arg Leu Leu Leu  
 501 TTG CTC AAG CAG ATT TAT CGC CAG CAG CTC  
 158 ◀ Gln Glu Leu Leu Asn Ile Ala Leu Leu Glu  
 531 CGA ATA GCG CCC TTC CCC TTG CCC GGC GTT  
 148 ◀ Ser Tyr Arg Gly Glu Gly Gln Gly Ala Asn  
 561 AAT GAT TTG CCC AAA CAG GTC GCT GAA ATG  
 138 ◀ Ile Ile Gln Gly Phe Leu Asp Ser Phe His  
 591 CGG CTG GTG CGC TTC ATC CGG GCG AAA GAA  
 128 ◀ Pro Gln His Ala Glu Asp Pro Arg Phe Phe  
 621 CCC CGT ATT GGC AAA TAT TGA CGG CCA GTT  
 118 ◀ Gly Thr Asn Ala Phe Ile Ser Pro Trp Asn  
 651 AAG CCA TTC ATG CCA GTA GGC GCG CGG ACC  
 108 ◀ Leu Trp Glu His Trp Tyr Ala Arg Pro Arg  
 681 AAA GTA AAC CCA CTG GTG ATA CCA TTC GCG  
 98 ◀ Phe Tyr Val Trp Gln His Tyr Trp Glu Arg  
 711 AGC CTC CGG ATG ACG ACC GTA GTG ATG AAT  
 88 ◀ Ala Glu Pro His Arg Gly Tyr His His Ile  
 741 CTC TCC TGG CGG GAA CAG CAA AAT ATC ACC  
 78 ◀ Glu Gly Pro Pro Phe Leu Leu Ile Asp Gly  
 771 CGG TCG GCA AAC AAA TTC TCG TCC CTG ATT  
 68 ◀ Pro Arg Cys Val Phe Glu Arg Gly Gln Asn

38/65

Figure 13b (cont'd)

801 TTT CAC CAC CCC CTG ACC GCG AAT GGT GAG  
58◀ Lys Val Val Gly Gln Gly Arg Ile Thr Leu

831 ATT GAG AAT ATA ACC TTT CAT TCC CAG CGG  
48◀ Asn Leu Ile Tyr Gly Lys Met Gly Leu Pro

861 TCG GTC GAT AAA AAA ATC GAG ATA ACC GTT  
38◀ Arg Asp Ile Phe Phe Asp Leu Tyr Gly Asn

891 GGC CTC AAT CGG CGT TAA ACC CGC CAC CAG  
28◀ Ala Glu Ile Pro Thr Leu Gly Ala Val Leu

921 ATG GGC ATT AAA CGA GTA TCC CGG CAG CAG  
18◀ His Ala Asn Phe Ser Tyr Gly Pro Leu Leu

951 GGG ATC ATT TTG CGC TTC AGC CAT ACTTTTC  
8◀ Pro Asp Asn Gln Ala Glu Ala Met

982 ATACTCCCGCCATTCAGAGAAGAAACCAATTGTCCATAT

---

1021 TGCATCAGACATTGCCGTCACCTGCGTCTTTTACTGGCTC

---

1060 TTCTCGCTAACCAAACCGGTAACCCCGCTTATTAAAAGC

---

1099 ATTCTGTAACAAAGCGGGACCAAAGCCATGACAAAACG

---

1138 CGTAACAAAAGTGTCTATAATCACGGCAGAAAAGTCCAC


---

1177 ATTGATTATTTGCACGGCGTCACACTTTGCTATGCCATA

---

1216 GCATTTTTTATCCATAAGATTAGCGGATCCTACCTGACGC

BamHI



SUBSTITUTE SHEET (RULE 26)

39/65

Figure 13b (cont'd)

1255 TTTTATCGCAACTCTCTACTGTTTCTCCATACCCGTTT  
                     NheI                    EcoRI            NcoI    BamHI  
 1294 TTTTGGGCTAGCAGGAGGAAT TCACC ATG GAT CCC  
   1►Met Asp Pro  
 1329 GTA ATC GTA GAA GAC ATA GAG CCA GGT ATT  
       4►Val Ile Val Glu Asp Ile Glu Pro Gly Ile  
 1359 TAT TAC GGA ATT TCG AAT GAG AAT TAC CAC  
       14►Tyr Tyr Gly Ile Ser Asn Glu Asn Tyr His  
 1389 GCG GGT CCC GGT ATC AGT AAG TCT CAG CTC  
       24►Ala Gly Pro Gly Ile Ser Lys Ser Gln Leu  
 1419 GAT GAC ATT GCT GAT ACT CCG GCA CTA TAT  
       34►Asp Asp Ile Ala Asp Thr Pro Ala Leu Tyr  
 1449 TTG TGG CGT AAA AAT GCC CCC GTG GAC ACC  
       44►Leu Trp Arg Lys Asn Ala Pro Val Asp Thr  
 1479 ACA AAG ACA AAA ACG CTC GAT TTA GGA ACT  
       54►Thr Lys Thr Lys Thr Leu Asp Leu Gly Thr  
 1509 GCT TTC CAC TGC CGG GTA CTT GAA CCG GAA  
       64►Ala Phe His Cys Arg Val Leu Glu Pro Glu  
                     EcoRI  
 1539 GAA TTC AGT AAC CGC TTT ATC GTA GCA CCT  
       74►Glu Phe Ser Asn Arg Phe Ile Val Ala Pro  
 1569 GAA TTT AAC CGC CGT ACA AAC GCC GGA AAA  
       84►Glu Phe Asn Arg Arg Thr Asn Ala Gly Lys  
 1599 GAA GAA GAG AAA GCG TTT CTG ATG GAA TGC  
       94►Glu Glu Glu Lys Ala Phe Leu Met Glu Cys  
 1629 GCA AGC ACA GGA AAA ACG GTT ATC ACT GCC  
       104►Ala Ser Thr Gly Lys Thr Val Ile Thr Ala

40/65

Figure 13b (cont'd)

1659	GAA	GAA	GGC	CGG	AAA	ATT	GAA	CTC	ATG	TAT
114▶	Glu	Glu	Gly	Arg	Lys	Ile	Glu	Leu	Met	Tyr
1689	CAA	AGC	GTT	ATG	GCT	TTG	CCG	CTG	GGG	CAA
124▶	Gln	Ser	Val	Met	Ala	Leu	Pro	Leu	Gly	Gln
1719	TGG	CTT	GTT	GAA	AGC	GCC	GGA	CAC	GCT	GAA
134▶	Trp	Leu	Val	Glu	Ser	Ala	Gly	His	Ala	Glu
1749	TCA	TCA	ATT	TAC	TGG	GAA	GAT	CCT	GAA	ACA
144▶	Ser	Ser	Ile	Tyr	Trp	Glu	Asp	Pro	Glu	Thr
1779	GGA	ATT	TTG	TGT	CGG	TGC	CGT	CCG	GAC	AAA
154▶	Gly	Ile	Leu	Cys	Arg	Cys	Arg	Pro	Asp	Lys
1809	ATT	ATC	CCT	GAA	TTT	CAC	TGG	ATC	ATG	GAC
164▶	Ile	Ile	Pro	Glu	Phe	His	Trp	Ile	Met	Asp
1839	GTG	AAA	ACT	ACG	GCG	GAT	ATT	CAA	CGA	TTC
174▶	Val	Lys	Thr	Thr	Ala	Asp	Ile	Gln	Arg	Phe
1869	AAA	ACC	GCT	TAT	TAC	GAC	TAC	CGC	TAT	CAC
184▶	Lys	Thr	Ala	Tyr	Tyr	Asp	Tyr	Arg	Tyr	His
1899	GTT	CAG	GAT	GCA	TTC	TAC	AGT	GAC	GGT	TAT
194▶	Val	Gln	Asp	Ala	Phe	Tyr	Ser	Asp	Gly	Tyr
1929	GAA	GCA	CAG	TTT	GGA	GTG	CAG	CCA	ACT	TTC
204▶	Glu	Ala	Gln	Phe	Gly	Val	Gln	Pro	Thr	Phe
1959	GTT	TTT	CTG	GTT	GCC	AGC	ACA	ACT	ATT	GAA
214▶	Val	Phe	Leu	Val	Ala	Ser	Thr	Thr	Ile	Glu
1989	TGC	GGA	CGT	TAT	CCG	GTT	GAA	ATT	TTC	ATG
224▶	Cys	Gly	Arg	Tyr	Pro	Val	Glu	Ile	Phe	Met
2019	ATG	GGC	GAA	GAA	GCA	AAA	CTG	GCA	GGT	CAA
234▶	Met	Gly	Glu	Glu	Ala	Lys	Leu	Ala	Gly	Gln

41/65

Figure 13b (cont'd)

2049 CAG GAA TAT CAC CGC AAT CTG CGA ACC CTG  
 244▶ Gln Glu Tyr His Arg Asn Leu Arg Thr Leu  
 2079 TCT GAC TGC CTG AAT ACC GAT GAA TGG CCA  
 254▶ Ser Asp Cys Leu Asn Thr Asp Glu Trp Pro  
 2109 GCT ATT AAG ACA TTA TCA CTG CCC CGC TGG  
 264▶ Ala Ile Lys Thr Leu Ser Leu Pro Arg Trp  
XhoI KpnI  
 2139 GCT AAG GAA TAT GCA AAT GAC TAGATCTCGAG  
 274▶ Ala Lys Glu Tyr Ala Asn Asp  
 2171 GTACCCGAGCACGTGTTGACAATTAATCATCGGCATAGT  
 2210 ATATCGGCATAGTATAATACGACAAGGTGAGGAACTAAA  
 NcoI  
 2249 CC ATG GCT AAG CAA CCA CCA ATC GCA AAA  
 1▶ Met Ala Lys Gln Pro Pro Ile Ala Lys  
 2278 GCC GAT CTG CAA AAA ACT CAG GGA AAC CGT  
 10▶ Ala Asp Leu Gln Lys Thr Gln Gly Asn Arg  
 2308 GCA CCA GCA GCA GTT AAA AAT AGC GAC GTG  
 20▶ Ala Pro Ala Ala Val Lys Asn Ser Asp Val  
 2338 ATT AGT TTT ATT AAC CAG CCA TCA ATG AAA  
 30▶ Ile Ser Phe Ile Asn Gln Pro Ser Met Lys  
 2368 GAG CAA CTG GCA GCA GCT CTT CCA CGC CAT  
 40▶ Glu Gln Leu Ala Ala Ala Leu Pro Arg His  
 2398 ATG ACG GCT GAA CGT ATG ATC CGT ATC GCC  
 50▶ Met Thr Ala Glu Arg Met Ile Arg Ile Ala  
 2428 ACC ACA GAA ATT CGT AAA GTT CCG GCG TTA  
 60▶ Thr Thr Glu Ile Arg Lys Val Pro Ala Leu

42/65

Figure 13b (cont'd)

2458	GGA	AAC	TGT	GAC	ACT	ATG	AGT	TTT	GTC	AGT
70▶	Gly	Asn	Cys	Asp	Thr	Met	Ser	Phe	Val	Ser
2488	GCG	ATC	GTA	CAG	TGT	TCA	CAG	CTC	GGA	CTT
80▶	Ala	Ile	Val	Gln	Cys	Ser	Gln	Leu	Gly	Leu
2518	GAG	CCA	GGT	AGC	GCC	CTC	GGT	CAT	GCA	TAT
90▶	Glu	Pro	Gly	Ser	Ala	Leu	Gly	His	Ala	Tyr
2548	TTA	CTG	CCT	TTT	GGT	AAT	AAA	AAC	GAA	AAG
100▶	Leu	Leu	Pro	Phe	Gly	Asn	Lys	Asn	Glu	Lys
2578	AGC	GGT	AAA	AAG	AAC	GTT	CAG	CTA	ATC	ATT
110▶	Ser	Gly	Lys	Lys	Asn	Val	Gln	Leu	Ile	Ile
2608	GGC	TAT	CGC	GGC	ATG	ATT	GAT	CTG	GCT	CGC
120▶	Gly	Tyr	Arg	Gly	Met	Ile	Asp	Leu	Ala	Arg
2638	CGT	TCT	GGT	CAA	ATC	GCC	AGC	CTG	TCA	GCC
130▶	Arg	Ser	Gly	Gln	Ile	Ala	Ser	Leu	Ser	Ala
2668	CGT	GTT	GTC	CGT	GAA	GGT	GAC	GAG	TTT	AGC
140▶	Arg	Val	Val	Arg	Glu	Gly	Asp	Glu	Phe	Ser
2698	TTC	GAA	TTT	GGC	CTT	GAT	GAA	AAG	TTA	ATA
150▶	Phe	Glu	Phe	Gly	Leu	Asp	Glu	Lys	Leu	Ile
2728	CAC	CGC	CCG	GGA	GAA	AAC	GAA	GAT	GCC	CCG
160▶	His	Arg	Pro	Gly	Glu	Asn	Glu	Asp	Ala	Pro
2758	GTT	ACC	CAC	GTC	TAT	GCT	GTC	GCA	AGA	CTG
170▶	Val	Thr	His	Val	Tyr	Ala	Val	Ala	Arg	Leu
2788	AAA	GAC	GGA	GGT	ACT	CAG	TTT	GAA	GTT	ATG
180▶	Lys	Asp	Gly	Gly	Thr	Gln	Phe	Glu	Val	Met
2818	ACG	CGC	AAA	CAG	ATT	GAG	CTG	GTG	CGC	AGC
190▶	Thr	Arg	Lys	Gln	Ile	Glu	Leu	Val	Arg	Ser

SUBSTITUTE SHEET (RULE 26)



43/65

Figure 13b (cont'd)

2848 CTG AGT AAA GCT GGT AAT AAC GGG CCG TGG  
200► Leu Ser Lys Ala Gly Asn Asn Gly Pro Trp

2878 GTA ACT CAC TGG GAA GAA ATG GCA AAG AAA  
210► Val Thr His Trp Glu Glu Met Ala Lys Lys

2908 ACG GCT ATT CGT CGC CTG TTC AAA TAT TTG  
220► Thr Ala Ile Arg Arg Leu Phe Lys Tyr Leu

2938 CCC GTA TCA ATT GAG ATC CAG CGT GCA GTA  
230► Pro Val Ser Ile Glu Ile Gln Arg Ala Val

2968 TCA ATG GAT GAA AAG GAA CCA CTG ACA ATC  
240► Ser Met Asp Glu Lys Glu Pro Leu Thr Ile

2998 GAT CCT GCA GAT TCC TCT GTA TTA ACC GGG  
250► Asp Pro Ala Asp Ser Ser Val Leu Thr Gly

3028 GAA TAC AGT GTA ATC GAT AAT TCA GAG GAA  
260► Glu Tyr Ser Val Ile Asp Asn Ser Glu Glu

BglII HindIII

3058 TAG ATCTAAGCTTCCTGCTGAACATCAAAGGCAAGAAA  
270► . . .

3096 ACATCTGTTGTCAAAGACAGCATCCTTGAACAAGGACAA

3135 TTAACAGTTAACAAATAAAAACGCAAAAGAAAATGCCGA

3174 TATCCTATTGGCATTTCCTTTTATTTCTTATCAACATAA

XhoI

3213 AGGTGAATCCCATACCTCGAGCTTCACGCTGCCGCAAGC

3252 ACTCAGGGCGCAAGGGCTGCTAAAAGGAAGCGGAACACG

3291 TAGAAAGCCAGTCCGCAGAAACGGTGCTGACCCCGGATG

3330 AATGTCAGCTACTGGGCTATCTGGACAAGGGAAAACGCA

3369 AGCGCAAAGAGAAAGCAGGTAGCTTGCAGTGGGGCTTACA

Figure 13b (cont'd)

3408 TGGCGATAGCTAGACTGGGCGGTTTATGGACAGCAAGC

3447 GAACCGGAATTGCCAGCTGGGGCGCCCTCTGGTAAGGTT

3486 GGGAAGCCCTGCAAAGTAAACTGGATGGCTTTCTTGCCG

BglII

3525 CCAAGGATCTGATGGCGCAGGGGATCAAGATCTGATCAA

3564 GAGACAGGATGAGGATCGTTTCGC ATG GAT ATT

1►Met Asp Ile

3597 AAT ACT GAA ACT GAG ATC AAG CAA AAG CAT

4►Asn Thr Glu Thr Glu Ile Lys Gln Lys His

3627 TCA CTA ACC CCC TTT CCT GTT TTC CTA ATC

14►Ser Leu Thr Pro Phe Pro Val Phe Leu Ile

3657 AGC CCG GCA TTT CGC GGG CGA TAT TTT CAC

24►Ser Pro Ala Phe Arg Gly Arg Tyr Phe His

3687 AGC TAT TTC AGG AGT TCA GCC ATG AAC GCT

34►Ser Tyr Phe Arg Ser Ser Ala Met Asn Ala

3717 TAT TAC ATT CAG GAT CGT CTT GAG GCT CAG

44►Tyr Tyr Ile Gln Asp Arg Leu Glu Ala Gln

3747 AGC TGG GCG CGT CAC TAC CAG CAG CTC GCC

54►Ser Trp Ala Arg His Tyr Gln Gln Leu Ala

3777 CGT GAA GAG AAA GAG GCA GAA CTG GCA GAC

64►Arg Glu Glu Lys Glu Ala Glu Leu Ala Asp

3807 GAC ATG GAA AAA GGC CTG CCC CAG CAC CTG

74►Asp Met Glu Lys Gly Leu Pro Gln His Leu

3837 TTT GAA TCG CTA TGC ATC GAT CAT TTG CAA

84►Phe Glu Ser Leu Cys Ile Asp His Leu Gln

3867 CGC CAC GGG GCC AGC AAA AAA TCC ATT ACC

94►Arg His Gly Ala Ser Lys Lys Ser Ile Thr

SUBSTITUTE SHEET (RULE 26)

45/65

Figure 13b (cont'd)

3897 CGT GCG TTT GAT GAC GAT GTT GAG TTT CAG  
 104▶ Arg Ala Phe Asp Asp Asp Val Glu Phe Gln  
 3927 GAG CGC ATG GCA GAA CAC ATC CGG TAC ATG  
 114▶ Glu Arg Met Ala Glu His Ile Arg Tyr Met  
 3957 GTT GAA ACC ATT GCT CAC CAC CAG GTT GAT  
 124▶ Val Glu Thr Ile Ala His His Gln Val Asp  
 HindIII  
 3987 ATT GAT TCA GAG GTA TAA AACGAGTAGA AGCT  
 134▶ Ile Asp Ser Glu Val ...  
 4019 TGGCTGTTTTGGCGGATGAGAGAAGATTTTCAGCCTGAT  
 4058 ACAGATTAAATCAGAACGCAGAAAGCGGTCTGATAAAACA  
 4097 GAATTTGCCTGGCGGCAGTAGCGCGGTGGTCCACCTGA  
 4136 CCCCATGCCGAACCTCAGAAGTGAAACGCCGTAGCGCCGA  
 4175 TGGTAGTGTGGGGTCTCCCCATGCGAGAGTAGGGAACTG  
 4214 CCAGGCATCAAATAAAACGAAAGGCTCAGTCGAAAGACT  
 4253 GGGCCTTTTCGTTTTATCTGTTGTTTGTCGGTGAACGCTC  
 4292 TCCTGAGTAGGACAAATCCGCCGGGAGCGGATTTGAACG  
 4331 TTGCGAAGCAACGGCCCCGGAGGGTGGCGGGCAGGACGCC  
 4370 CGCCATAAACTGCCAGGCATCAAATTAAGCAGAAGGCCA  
 4409 TCCTGACGGATGGCCTTTTTTGCGTTTCTACAAACTCTTT  
 4448 TGTTTATTTTTTCTAAATACATTCAAATATGTATCCGCTC  
 4487 ATGAGACAATAACCCTGATAAATGCTTCAATAATATTGA  
 4526 AAAAGGAAGAGT ATG AGT ATT CAA CAT TTC  
 1▶ Met Ser Ile Gln His Phe

Figure 13b (cont'd)

4556	CGT	GTC	GCC	CTT	ATT	CCC	TTT	TTT	GCG	GCA
7▶	Arg	Val	Ala	Leu	Ile	Pro	Phe	Phe	Ala	Ala
4586	TTT	TGC	CTT	CCT	GTT	TTT	GCT	CAC	CCA	GAA
17▶	Phe	Cys	Leu	Pro	Val	Phe	Ala	His	Pro	Glu
4616	ACG	CTG	GTG	AAA	GTA	AAA	GAT	GCT	GAA	GAT
27▶	Thr	Leu	Val	Lys	Val	Lys	Asp	Ala	Glu	Asp
4646	CAG	TTG	GGT	GCA	CGA	GTG	GGT	TAC	ATC	GAA
37▶	Gln	Leu	Gly	Ala	Arg	Val	Gly	Tyr	Ile	Glu
4676	CTG	GAT	CTC	AAC	AGC	GGT	AAG	ATC	CTT	GAG
47▶	Leu	Asp	Leu	Asn	Ser	Gly	Lys	Ile	Leu	Glu
4706	AGT	TTT	CGC	CCC	GAA	GAA	CGT	TTT	CCA	ATG
57▶	Ser	Phe	Arg	Pro	Glu	Glu	Arg	Phe	Pro	Met
4736	ATG	AGC	ACT	TTT	AAA	GTT	CTG	CTA	TGT	GGC
67▶	Met	Ser	Thr	Phe	Lys	Val	Leu	Leu	Cys	Gly
4766	GCG	GTA	TTA	TCC	CGT	GTT	GAC	GCC	GGG	CAA
77▶	Ala	Val	Leu	Ser	Arg	Val	Asp	Ala	Gly	Gln
4796	GAG	CAA	CTC	GGT	CGC	CGC	ATA	CAC	TAT	TCT
87▶	Glu	Gln	Leu	Gly	Arg	Arg	Ile	His	Tyr	Ser
						Scal				
4826	CAG	AAT	GAC	TTG	GTT	GAG	TAC	TCA	CCA	GTC
97▶	Gln	Asn	Asp	Leu	Val	Glu	Tyr	Ser	Pro	Val
4856	ACA	GAA	AAG	CAT	CTT	ACG	GAT	GGC	ATG	ACA
107▶	Thr	Glu	Lys	His	Leu	Thr	Asp	Gly	Met	Thr
4886	GTA	AGA	GAA	TTA	TGC	AGT	GCT	GCC	ATA	ACC
117▶	Val	Arg	Glu	Leu	Cys	Ser	Ala	Ala	Ile	Thr
4916	ATG	AGT	GAT	AAC	ACT	GCG	GCC	AAC	TTA	CTT
127▶	Met	Ser	Asp	Asn	Thr	Ala	Ala	Asn	Leu	Leu

47/65

Figure 13b (cont'd)

4946	CTG	ACA	ACG	ATC	GGA	GGA	CCG	AAG	GAG	CTA
137▶	Leu	Thr	Thr	Ile	Gly	Gly	Pro	Lys	Glu	Leu
4976	ACC	GCT	TTT	TTG	CAC	AAC	ATG	GGG	GAT	CAT
147▶	Thr	Ala	Phe	Leu	His	Asn	Met	Gly	Asp	His
5006	GTA	ACT	CGC	CTT	GAT	CGT	TGG	GAA	CCG	GAG
157▶	Val	Thr	Arg	Leu	Asp	Arg	Trp	Glu	Pro	Glu
5036	CTG	AAT	GAA	GCC	ATA	CCA	AAC	GAC	GAG	CGT
167▶	Leu	Asn	Glu	Ala	Ile	Pro	Asn	Asp	Glu	Arg
5066	GAC	ACC	ACG	ATG	CCT	GTA	GCA	ATG	GCA	ACA
177▶	Asp	Thr	Thr	Met	Pro	Val	Ala	Met	Ala	Thr
5096	ACG	TTG	CGC	AAA	CTA	TTA	ACT	GGC	GAA	CTA
187▶	Thr	Leu	Arg	Lys	Leu	Leu	Thr	Gly	Glu	Leu
5126	CTT	ACT	CTA	GCT	TCC	CGG	CAA	CAA	TTA	ATA
197▶	Leu	Thr	Leu	Ala	Ser	Arg	Gln	Gln	Leu	Ile
5156	GAC	TGG	ATG	GAG	GCG	GAT	AAA	GTT	GCA	GGA
207▶	Asp	Trp	Met	Glu	Ala	Asp	Lys	Val	Ala	Gly
5186	CCA	CTT	CTG	CGC	TCG	GCC	CTT	CCG	GCT	GGC
217▶	Pro	Leu	Leu	Arg	Ser	Ala	Leu	Pro	Ala	Gly
5216	TGG	TTT	ATT	GCT	GAT	AAA	TCT	GGA	GCC	GGT
227▶	Trp	Phe	Ile	Ala	Asp	Lys	Ser	Gly	Ala	Gly
5246	GAG	CGT	GGG	TCT	CGC	GGT	ATC	ATT	GCA	GCA
237▶	Glu	Arg	Gly	Ser	Arg	Gly	Ile	Ile	Ala	Ala
5276	CTG	GGG	CCA	GAT	GGT	AAG	CCC	TCC	CGT	ATC
247▶	Leu	Gly	Pro	Asp	Gly	Lys	Pro	Ser	Arg	Ile
5306	GTA	GTT	ATC	TAC	ACG	ACG	GGG	AGT	CAG	GCA
257▶	Val	Val	Ile	Tyr	Thr	Thr	Gly	Ser	Gln	Ala

Figure 13b (cont'd)

5336 ACT ATG GAT GAA CGA AAT AGA CAG ATC GCT  
267▶ Thr Met Asp Glu Arg Asn Arg Gln Ile Ala

5366 GAG ATA GGT GCC TCA CTG ATT AAG CAT TGG  
277▶ Glu Ile Gly Ala Ser Leu Ile Lys His Trp

5396 TAA CTGTCAGACCAAGTTTACTCATATATACTTTAGAT  
287▶ . . .

5434 TGATTTACGCGCCCTGTAGCGGCGCATTAAAGCGCGGCGG  
5473 GTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCA  
5512 GCGCCCTAGCGCCCGCTCCTTTTCGCTTTCTTCCCTTCCT  
5551 TTCTCGCCACGTTTCGCCGGCTTTCCCCGTCAAGCTCTAA  
5590 ATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTAC  
5629 GGCACCTCGACCCCCAAAAAACTTGATTTGGGTGATGGTT  
5668 CACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTTCGCC  
5707 CTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCT  
5746 TGTTCCAAACCTTGAACAACACTCAACCCTATCTCGGGCT  
5785 ATTCTTTTGATTTTATAAGGGATTTTGCCGATTTTCGGCCT  
5824 ATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAACG  
5863 CGAATTTTAACAAAATATTAACGTTTACAATTTAAAAGG  
5902 ATCTAGGTGAAGATCCTTTTTTGATAATCTCATGACCAAA  
5941 ATCCCTTAACGTGAGTTTTTCGTTCCACTGAGCGTCAGAC  
5980 CCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTT  
6019 TTTCTGCGCGTAATCTGCTGCTTGCAAACAAAAAAACCA  
6058 CCGCTACCAGCGGTGGTTTGTTTGCCGGATCAAGAGCTA  
6097 CCAACTCTTTTTTCCGAAGGTAAGTGGCTTCAGCAGAGCG

49/65

Figure 13b (cont'd)

6136 CAGATACCAAATACTGTCCTTCTAGTGTAGCCGTAGTTA  
6175 GGCCACCACCTTCAAGAACTCTGTAGCACCGCCTACATAC  
6214 CTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGT  
6253 GGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGA  
6292 TAGTTACCGGATAAGGCGCAGCGGTCTGGGCTGAACGGGG  
6331 GGTTCGTGCACACAGCCCAGCTTGGAGCGAACGACCTAC  
6370 ACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAGC  
6409 GCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCG  
6448 GTAAGCGGCAGGGTTCGGAACAGGAGAGCGCACGAGGGAG  
6487 CTTCCAGGGGGGAAACGCCTGGGTATCTTTATAGTCCTGTC  
6526 GGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGA  
6565 TGCTCGTCAGGGGGGGCGGAGCCTATGGAAAAACGCCAGC  
6604 AACGCGGCCTTTTTACGGTTCCTGGCCTTTTGCTGGCCT  
6643 TTTGCTCACATGTTCTTTCTGCGTTATCCCCTGATTCT  
6682 GTGGATAACCGTATTACCGCCTTTGAGTGAGCTGATACC  
6721 GCTCGCCGCAGCCGAACGACCGAGCGCAGCGAGTCAGTG  
6760 AGCGAGGAAGCGGAAGAGCGCCTGATGCGGTATTTTCTC  
6799 CTTACGCATCTGTGCGGTATTTTCACACCGCATAGGGTCA  
6838 TGGCTGCGCCCCGACACCCGCCAACACCCGCTGACGCGC  
6877 CCTGACGGGCTTGTCTGCTCCCGGCATCCGCTTACAGAC  
6916 AAGCTGTGACCGTCTCCGGGAGCTGCATGTGTCAGAGGT  
6955 TTTCACCGTCATCACCGAAACGCGCGAGGCAGCAAGGAG

50/65


Figure 13b (cont'd)

6994 ATGGCGCCCAACAGTCCCCCGGCCACGGGGCCTGCCACC  
7033 ATACCCACGCCGAAACAAGCGCTCATGAGCCCGAAGTGG  
7072 CGAGCCCGATCTTCCCCATCGGGTGATGTCGGCGATATAG  
7111 GCGCCAGCAACCGCACCTGTGGCGCCGGTGATGCCGGCC  
7150 ACGATGCGTCCGGCGTAGAGGATCTGCTCATGTTTGACA  
7189 GCTTATC

SUBSTITUTE SHEET (RULE 26)



Figure 14 a



52:65

Figure 14b

## NsiI

1 ATCGATGCATAATGTGCCTGTCAAATGGACGAAGCAGGG  
 40 ATTCTGCAAACCCTATGCTACTCCGTCAAGCCGTCAATT  
 79 GTCTGATTTCGTTACCAA TTA TGA CAA CTT GAC  
 293◀••• Ser Leu Lys Val  
 111 GGC TAC ATC ATT CAC TTT TTC TTC ACA ACC  
 288◀Ala Val Asp Asn Val Lys Glu Glu Cys Gly  
 141 GGC ACG GAA CTC GCT CGG GCT GGC CCC GGT  
 278◀Ala Arg Phe Glu Ser Pro Ser Ala Gly Thr  
 171 GCA TTT TTT AAA TAC CCG CGA GAA ATA GAG  
 268◀Cys Lys Lys Phe Val Arg Ser Phe Tyr Leu  
 201 TTG ATC GTC AAA ACC AAC ATT GCG ACC GAC  
 258◀Gln Asp Asp Phe Gly Val Asn Arg Gly Val  
 231 GGT GGC GAT AGG CAT CCG GGT GGT GCT CAA  
 248◀Thr Ala Ile Pro Met Arg Thr Thr Ser Leu  
 261 AAG CAG CTT CGC CTG GCT GAT ACG TTG GTC  
 238◀Leu Leu Lys Ala Gln Ser Ile Arg Gln Asp  
 291 CTC GCG CCA GCT TAA GAC GCT AAT CCC TAA  
 228◀Glu Arg Trp Ser Leu Val Ser Ile Gly Leu  
 321 CTG CTG GCG GAA AAG ATG TGA CAG ACG CGA  
 218◀Gln Gln Arg Phe Leu His Ser Leu Arg Ser  
 351 CGG CGA CAA GCA AAC ATG CTG TGC GAC GCT  
 208◀Pro Ser Leu Cys Val His Gln Ala Val Ser  
 EcoRV  
 381 GGC GAT ATC AAA ATT GCT GTC TGC CAG GTG  
 198◀Ala Ile Asp Phe Asn Ser Asp Ala Leu His

SUBSTITUTE SHEET (RULE 26)

53/65

Figure 14b (cont'd)

411	ATC	GCT	GAT	GTA	CTG	ACA	AGC	CTC	GCG	TAC
188	◀Asp	Ser	Ile	Tyr	Gln	Cys	Ala	Glu	Arg	Val
441	CCG	ATT	ATC	CAT	CGG	TGG	ATG	GAG	CGA	CTC
178	◀Arg	Asn	Asp	Met	Pro	Pro	His	Leu	Ser	Glu
471	GTT	AAT	CGC	TTC	CAT	GCG	CCG	CAG	TAA	CAA
168	◀Asn	Ile	Ala	Glu	Met	Arg	Arg	Leu	Leu	Leu
501	TTG	CTC	AAG	CAG	ATT	TAT	CGC	CAG	CAG	CTC
158	◀Gln	Glu	Leu	Leu	Asn	Ile	Ala	Leu	Leu	Glu
531	CGA	ATA	GCG	CCC	TTC	CCC	TTG	CCC	GGC	GTT
148	◀Ser	Tyr	Arg	Gly	Glu	Gly	Gln	Gly	Ala	Asn
561	AAT	GAT	TTG	CCC	AAA	CAG	GTC	GCT	GAA	ATG
138	◀Ile	Ile	Gln	Gly	Phe	Leu	Asp	Ser	Phe	His
591	CGG	CTG	GTG	CGC	TTC	ATC	CGG	GCG	AAA	GAA
128	◀Pro	Gln	His	Ala	Glu	Asp	Pro	Arg	Phe	Phe
621	CCC	CGT	ATT	GGC	AAA	TAT	TGA	CGG	CCA	GTT
118	◀Gly	Thr	Asn	Ala	Phe	Ile	Ser	Pro	Trp	Asn
651	AAG	CCA	TTC	ATG	CCA	GTA	GGC	GCG	CGG	ACG
108	◀Leu	Trp	Glu	His	Trp	Tyr	Ala	Arg	Pro	Arg
681	AAA	GTA	AAC	CCA	CTG	GTG	ATA	CCA	TTC	GCG
98	◀Phe	Tyr	Val	Trp	Gln	His	Tyr	Trp	Glu	Arg
711	AGC	CTC	CGG	ATG	ACG	ACC	GTA	GTG	ATG	AAT
88	◀Ala	Glu	Pro	His	Arg	Gly	Tyr	His	His	Ile
741	CTC	TCC	TGG	CGG	GAA	CAG	CAA	AAT	ATC	ACC
78	◀Glu	Gly	Pro	Pro	Phe	Leu	Leu	Ile	Asp	Gly
771	CGG	TCG	GCA	AAC	AAA	TTC	TCG	TCC	CTG	ATT
68	◀Pro	Arg	Cys	Val	Phe	Glu	Arg	Gly	Gln	Asn

54/65

Figure 14b (cont'd)

801 TTT CAC CAC CCC CTG ACC GCG AAT GGT GAG  
 58◀ Lys Val Val Gly Gln Gly Arg Ile Thr Leu  
 831 ATT GAG AAT ATA ACC TTT CAT TCC CAG CGG  
 48◀ Asn Leu Ile Tyr Gly Lys Met Gly Leu Pro  
 861 TCG GTC GAT AAA AAA ATC GAG ATA ACC GTT  
 38◀ Arg Asp Ile Phe Phe Asp Leu Tyr Gly Asn  
 891 GGC CTC AAT CGG CGT TAA ACC CGC CAC CAG  
 28◀ Ala Glu Ile Pro Thr Leu Gly Ala Val Leu  
 921 ATG GGC ATT AAA CGA GTA TCC CGG CAG CAG  
 18◀ His Ala Asn Phe Ser Tyr Gly Pro Leu Leu  
 951 GGG ATC ATT TTG CGC TTC AGC CAT ACTTTTC  
 8◀ Pro Asp Asn Gln Ala Glu Ala Met  
 982 ATACTCCCGCCATTCAGAGAAGAAACCAATTGTCCATAT  
 1021 TGCATCAGACATTGCCGTCCTGCGTCTTTTACTGGCTC  
 1060 TTCTCGCTAACCAAACCGGTAACCCCGCTTATTAAAAGC  
 1099 ATTCTGTAACAAAGCGGGACCAAAGCCATGACAAAAACG  
 1138 CGTAACAAAAGTGTCTATAATCACGGCAGAAAAGTCCAC  
 1177 ATTGATTATTTGCACGGCGTCACACTTTTGCTATGCCATA  
 BamHI  
 1216 GCATTTTTATCCATAAGATTAGCGGATCCTACCTGACGC  
 1255 TTTTATCGCAACTCTCTACTGTTTCTCCATACCCGTTT  
 NheI EcoRI  
 1294 TTTTGGGCTAGCAGGAGGAATTCACC ATG ACA CCG  
 1▶ Met Thr Pro  
 PstI  
 1329 GAC ATT ATC CTG CAG CGT ACC GGG ATC GAT

55/65

Figure 14b (cont'd)

4▶	Asp	Ile	Ile	Leu	Gln	Arg	Thr	Gly	Ile	Asp
1359	GTG	AGA	GCT	GTC	GAA	CAG	GGG	GAT	GAT	GCG
14▶	Val	Arg	Ala	Val	Glu	Gln	Gly	Asp	Asp	Ala
1389	TGG	CAC	AAA	TTA	CGG	CTC	GGC	GTC	ATC	ACC
24▶	Trp	His	Lys	Leu	Arg	Leu	Gly	Val	Ile	Thr
1419	GCT	TCA	GAA	GTT	CAC	AAC	GTG	ATA	GCA	AAA
34▶	Ala	Ser	Glu	Val	His	Asn	Val	Ile	Ala	Lys
1449	CCC	CGC	TCC	GGA	AAG	AAG	TGG	CCT	GAC	ATG
44▶	Pro	Arg	Ser	Gly	Lys	Lys	Trp	Pro	Asp	Met
1479	AAA	ATG	TCC	TAC	TTC	CAC	ACC	CTG	CTT	GCT
54▶	Lys	Met	Ser	Tyr	Phe	His	Thr	Leu	Leu	Ala
1509	GAG	GTT	TGC	ACC	GGT	GTG	GCT	CCG	GAA	GTT
64▶	Glu	Val	Cys	Thr	Gly	Val	Ala	Pro	Glu	Val
1539	AAC	GCT	AAA	GCA	CTG	GCC	TGG	GGA	AAA	CAG
74▶	Asn	Ala	Lys	Ala	Leu	Ala	Trp	Gly	Lys	Gln
										EcoRI
1569	TAC	GAG	AAC	GAC	GCC	AGA	ACC	CTG	TTT	GAA
84▶	Tyr	Glu	Asn	Asp	Ala	Arg	Thr	Leu	Phe	Glu
1599	TTC	ACT	TCC	GGC	GTG	AAT	GTT	ACT	GAA	TCC
94▶	Phe	Thr	Ser	Gly	Val	Asn	Val	Thr	Glu	Ser
1629	CCG	ATC	ATC	TAT	CGC	GAC	GAA	AGT	ATG	CGT
104▶	Pro	Ile	Ile	Tyr	Arg	Asp	Glu	Ser	Met	Arg
1659	ACC	GCC	TGC	TCT	CCC	GAT	GGT	TTA	TGC	AGT
114▶	Thr	Ala	Cys	Ser	Pro	Asp	Gly	Leu	Cys	Ser
1689	GAC	GGC	AAC	GGC	CTT	GAA	CTG	AAA	TGC	CCG
124▶	Asp	Gly	Asn	Gly	Leu	Glu	Leu	Lys	Cys	Pro

56/65

Figure 14b (cont'd)

1719 TTT ACC TCC CGG GAT TTC ATG AAG TTC CGG  
 134► Phe Thr Ser Arg Asp Phe Met Lys Phe Arg  
 1749 CTC GGT GGT TTC GAG GCC ATA AAG TCA GCT  
 144► Leu Gly Gly Phe Glu Ala Ile Lys Ser Ala  
 1779 TAC ATG GCC CAG GTG CAG TAC AGC ATG TGG  
 154► Tyr Met Ala Gln Val Gln Tyr Ser Met Trp  
 1809 GTG ACG CGA AAA AAT GCC TGG TAC TTT GCC  
 164► Val Thr Arg Lys Asn Ala Trp Tyr Phe Ala  
 1839 AAC TAT GAC CCG CGT ATG AAG CGT GAA GGC  
 174► Asn Tyr Asp Pro Arg Met Lys Arg Glu Gly  
 1869 CTG CAT TAT GTC GTG ATT GAG CGG GAT GAA  
 184► Leu His Tyr Val Val Ile Glu Arg Asp Glu  
 1899 AAG TAC ATG GCG AGT TTT GAC GAG ATC GTG  
 194► Lys Tyr Met Ala Ser Phe Asp Glu Ile Val  
 1929 CCG GAG TTC ATC GAA AAA ATG GAC GAG GCA  
 204► Pro Glu Phe Ile Glu Lys Met Asp Glu Ala  
 1959 CTG GCT GAA ATT GGT TTT GTA TTT GGG GAG  
 214► Leu Ala Glu Ile Gly Phe Val Phe Gly Glu  
 KpnI  
 1989 CAA TGG CGA TAGATCCGGTACCCGAGCACGTGTTGA  
 224► Gln Trp Arg . . .  
 2025 CAATTAATCATCGGCATAGTATATCGGCATAGTATAATA  
 2064 CGACAAGGTGAGGAACTAAACC ATG AGT ACT GCA  
 1► Met Ser Thr Ala  
 2098 CTC GCA ACG CTG GCT GGG AAG CTG GCT GAA  
 5► Leu Ala Thr Leu Ala Gly Lys Leu Ala Glu

SUBSTITUTE SHEET (RULE 26)

57/65

Figure 14b (cont'd)

	Sall									
2128	CGT	GTC	GGC	ATG	GAT	TCT	GTC	GAC	CCA	CAG
15▶	Arg	Val	Gly	Met	Asp	Ser	Val	Asp	Pro	Gln
2158	GAA	CTG	ATC	ACC	ACT	CTT	CGC	CAG	ACG	GCA
25▶	Glu	Leu	Ile	Thr	Thr	Leu	Arg	Gln	Thr	Ala
2188	TTT	AAA	GGT	GAT	GCC	AGC	GAT	GCG	CAG	TTC
35▶	Phe	Lys	Gly	Asp	Ala	Ser	Asp	Ala	Gln	Phe
2218	ATC	GCA	TTA	CTG	ATC	GTT	GCC	AAC	CAG	TAC
45▶	Ile	Ala	Leu	Leu	Ile	Val	Ala	Asn	Gln	Tyr
2248	GGC	CTT	AAT	CCG	TGG	ACG	AAA	GAA	ATT	TAC
55▶	Gly	Leu	Asn	Pro	Trp	Thr	Lys	Glu	Ile	Tyr
2278	GCC	TTT	CCT	GAT	AAG	CAG	AAT	GGC	ATC	GTT
65▶	Ala	Phe	Pro	Asp	Lys	Gln	Asn	Gly	Ile	Val
2308	CCG	GTG	GTG	GGC	GTT	GAT	GGC	TGG	TCC	CGC
75▶	Pro	Val	Val	Gly	Val	Asp	Gly	Trp	Ser	Arg
2338	ATC	ATC	AAT	GAA	AAC	CAG	CAG	TTT	GAT	GGC
85▶	Ile	Ile	Asn	Glu	Asn	Gln	Gln	Phe	Asp	Gly
2368	ATG	GAC	TTT	GAG	CAG	GAC	AAT	GAA	TCC	TGT
95▶	Met	Asp	Phe	Glu	Gln	Asp	Asn	Glu	Ser	Cys
2398	ACA	TGC	CGG	ATT	TAC	CGC	AAG	GAC	CGT	AAT
105▶	Thr	Cys	Arg	Ile	Tyr	Arg	Lys	Asp	Arg	Asn
2428	CAT	CCG	ATC	TGC	GTT	ACC	GAA	TGG	ATG	GAT
115▶	His	Pro	Ile	Cys	Val	Thr	Glu	Trp	Met	Asp
2458	GAA	TGC	CGC	CGC	GAA	CCA	TTC	AAA	ACT	CGC
125▶	Glu	Cys	Arg	Arg	Glu	Pro	Phe	Lys	Thr	Arg
2488	GAA	GGC	AGA	GAA	ATC	ACG	GGG	CCG	TGG	CAG
135▶	Glu	Gly	Arg	Glu	Ile	Thr	Gly	Pro	Trp	Gln

58/65

Figure 14b (cont'd)

2518 TCG CAT CCC AAA CGG ATG TTA CGT CAT AAA  
 145► Ser His Pro Lys Arg Met Leu Arg His Lys  
 2548 GCC ATG ATT CAG TGT GCC CGT CTG GCC TTC  
 155► Ala Met Ile Gln Cys Ala Arg Leu Ala Phe  
 2578 GGA TTT GCT GGT ATC TAT GAC AAG GAT GAA  
 165► Gly Phe Ala Gly Ile Tyr Asp Lys Asp Glu  
 2608 GCC GAG CGC ATT GTC GAA AAT ACT GCA TAC  
 175► Ala Glu Arg Ile Val Glu Asn Thr Ala Tyr  
 PstI  
 2638 ACT GCA GAA CGT CAG CCG GAA CGC GAC ATC  
 185► Thr Ala Glu Arg Gln Pro Glu Arg Asp Ile  
 2668 ACT CCG GTT AAC GAT GAA ACC ATG CAG GAG  
 195► Thr Pro Val Asn Asp Glu Thr Met Gln Glu  
 2698 ATT AAC ACT CTG CTG ATC GCC CTG GAT AAA  
 205► Ile Asn Thr Leu Leu Ile Ala Leu Asp Lys  
 2728 ACA TGG GAT GAC GAC TTA TTG CCG CTC TGT  
 215► Thr Trp Asp Asp Asp Leu Leu Pro Leu Cys  
 2758 TCC CAG ATA TTT CGC CGC GAC ATT CGT GCA  
 225► Ser Gln Ile Phe Arg Arg Asp Ile Arg Ala  
 2788 TCG TCA GAA CTG ACA CAG GCC GAA GCA GTA  
 235► Ser Ser Glu Leu Thr Gln Ala Glu Ala Val  
 2818 AAA GCT CTT GGA TTC CTG AAA CAG AAA GCC  
 245► Lys Ala Leu Gly Phe Leu Lys Gln Lys Ala  
 BglII XhoI  
 2848 GCA GAG CAG AAG GTG GCA GCA TAGATCTCGAG  
 255► Ala Glu Gln Lys Val Ala Ala ...

SUBSTITUTE SHEET (RULE 26)



59/65

Figure 14b (cont'd)

## HindIII

2880 AAGCTTCCTGCTGAACATCAAAGGCAAGAAAACATCTGT  
 2919 TGTCAAAGACAGCATCCTTGAACAAGGACAATTAACAGT  
 2958 TAACAAATAAAAAACGCAAAAGAAAATGCCGATATCCTAT  
 2997 TGGCATTTCCTTTTATTTCTTATCAACATAAAGGTGAAT

## XhoI

3036 CCCATACCTCGAGCTTCACGCTGCCGCAAGCACTCAGGG  
 3075 CGCAAGGGCTGCTAAAAGGAAGCGGAACACGTAGAAAGC  
 3114 CAGTCCGCAGAAACGGTGCTGACCCCGGATGAATGTCAG  
 3153 CTACTGGGCTATCTGGACAAGGGAAAACGCAAGCGCAAA  
 3192 GAGAAAGCAGGTAGCTTGCAGTGGGCTTACATGGCGATA  
 3231 GCTAGACTGGGCGGTTTTATGGACAGCAAGCGAACCGGA

## PvuII

3270 ATTGCCAGCTGGGGCGCCCTCTGGTAAGGTTGGGAAGCC  
 3309 CTGCAAAGTAAACTGGATGGCTTTCTTGCCGCCAAGGAT

## BglII

3348 CTGATGGCGCAGGGGATCAAGATCTGATCAAGAGACAGG  
 3387 ATGAGGATCGTTTCGC ATG GAT ATT AAT ACT

1►Met Asp Ile Asn Thr

3418 GAA ACT GAG ATC AAG CAA AAG CAT TCA CTA  
 6►Glu Thr Glu Ile Lys Gln Lys His Ser Leu

3448 ACC CCC TTT CCT GTT TTC CTA ATC AGC CCG  
 16►Thr Pro Phe Pro Val Phe Leu Ile Ser Pro

3478 GCA TTT CGC GGG CGA TAT TTT CAC AGC TAT  
 26►Ala Phe Arg Gly Arg Tyr Phe His Ser Tyr

3508 TTC AGG AGT TCA GCC ATG AAC GCT TAT TAC  
 36►Phe Arg Ser Ser Ala Met Asn Ala Tyr Tyr

60/65

Figure 14b (cont'd)

3538 ATT CAG GAT CGT CTT GAG GCT CAG AGC TGG  
 46► Ile Gln Asp Arg Leu Glu Ala Gln Ser Trp  
 3568 GCG CGT CAC TAC CAG CAG CTC GCC CGT GAA  
 56► Ala Arg His Tyr Gln Gln Leu Ala Arg Glu  
 3598 GAG AAA GAG GCA GAA CTG GCA GAC GAC ATG  
 66► Glu Lys Glu Ala Glu Leu Ala Asp Asp Met  
 3628 GAA AAA GGC CTG CCC CAG CAC CTG TTT GAA  
 76► Glu Lys Gly Leu Pro Gln His Leu Phe Glu  
 3658 TCG CTA TGC ATC GAT CAT TTG CAA CGC CAC  
 86► Ser Leu Cys Ile Asp His Leu Gln Arg His  
 3688 GGG GCC AGC AAA AAA TCC ATT ACC CGT GCG  
 96► Gly Ala Ser Lys Lys Ser Ile Thr Arg Ala  
 3718 TTT GAT GAC GAT GTT GAG TTT CAG GAG CGC  
 106► Phe Asp Asp Asp Val Glu Phe Gln Glu Arg  
 3748 ATG GCA GAA CAC ATC CGG TAC ATG GTT GAA  
 116► Met Ala Glu His Ile Arg Tyr Met Val Glu  
 3778 ACC ATT GCT CAC CAC CAG GTT GAT ATT GAT  
 126► Thr Ile Ala His His Gln Val Asp Ile Asp  
 HindIII  
 3808 TCA GAG GTA TAA AACGAGTAGA AGC TTG GCT  
 136► Ser Glu Val . . .  
 3839 GTT TTG GCG GAT GAG AGA AGA TTT TCA GCC  
 3869 TGA TACAGATTAAATCAGAACGCAGAAGCGGTCTGATA  
 3907 AAACAGAATTTGCCTGGCGGCAGTAGCGCGGTGGTCCCA  
 3946 CCTGACCCCATGCCGAAGTGAACGCCGTAGC  
 3985 GCCGATGGTAGTGTGGGGTCTCCCCATGCGAGAGTAGGG

SUBSTITUTE SHEET (RULE 26)

61/65

Figure 14b (cont'd)

4024 AACTGCCAGGCATCAAATAAAACGAAAGGCTCAGTCGAA  
 4063 AGACTGGGCCTTTTCGTTTTATCTGTTGTTTGTCGGTGAA  
 4102 CGCTCTCCTGAGTAGGACAAATCCGCCGGGAGCGGATTT  
 4141 GAACGTTGCGAAGCAACGGCCCCGGAGGGTGGCGGGCAGG  
 4180 ACGCCCGCCATAAACTGCCAGGCATCAAATTAAGCAGAA  
 4219 GGCCATCCTGACGGATGGCCTTTTTTGCGTTTCTACAAAC  
 4258 TCTTTTGTTTTATTTTTTCTAAATACATTCAAATATGTATC  
 4297 CGCTCATGAGACAATAACCCTGATAAATGCTTCAATAAT  
 4336 ATTGAAAAAGGAAGAGT ATG AGT ATT CAA CAT  
                                   1►Met Ser Ile Gln His  
 4368 TTC CGT GTC GCC CTT ATT CCC TTT TTT GCG  
       6►Phe Arg Val Ala Leu Ile Pro Phe Phe Ala  
 4398 GCA TTT TGC CTT CCT GTT TTT GCT CAC CCA  
       16►Ala Phe Cys Leu Pro Val Phe Ala His Pro  
 4428 GAA ACG CTG GTG AAA GTA AAA GAT GCT GAA  
       26►Glu Thr Leu Val Lys Val Lys Asp Ala Glu  
 4458 GAT CAG TTG GGT GCA CGA GTG GGT TAC ATC  
       36►Asp Gln Leu Gly Ala Arg Val Gly Tyr Ile  
 4488 GAA CTG GAT CTC AAC AGC GGT AAG ATC CTT  
       46►Glu Leu Asp Leu Asn Ser Gly Lys Ile Leu  
 4518 GAG AGT TTT CGC CCC GAA GAA CGT TTT CCA  
       56►Glu Ser Phe Arg Pro Glu Glu Arg Phe Pro  
 4548 ATG ATG AGC ACT TTT AAA GTT CTG CTA TGT  
       66►Met Met Ser Thr Phe Lys Val Leu Leu Cys

62/65

Figure 14b (cont'd)

4578	GGC	GCG	GTA	TTA	TCC	CGT	GTT	GAC	GCC	GGG
76▶	Gly	Ala	Val	Leu	Ser	Arg	Val	Asp	Ala	Gly
4608	CAA	GAG	CAA	CTC	GGT	CGC	CGC	ATA	CAC	TAT
86▶	Gln	Glu	Gln	Leu	Gly	Arg	Arg	Ile	His	Tyr
							Scal			
4638	TCT	CAG	AAT	GAC	TTG	GTT	GAG	TAC	TCA	CCA
96▶	Ser	Gln	Asn	Asp	Leu	Val	Glu	Tyr	Ser	Pro
4668	GTC	ACA	GAA	AAG	CAT	CTT	ACG	GAT	GGC	ATG
106▶	Val	Thr	Glu	Lys	His	Leu	Thr	Asp	Gly	Met
4698	ACA	GTA	AGA	GAA	TTA	TGC	AGT	GCT	GCC	ATA
116▶	Thr	Val	Arg	Glu	Leu	Cys	Ser	Ala	Ala	Ile
4728	ACC	ATG	AGT	GAT	AAC	ACT	GCG	GCC	AAC	TTA
126▶	Thr	Met	Ser	Asp	Asn	Thr	Ala	Ala	Asn	Leu
4758	CTT	CTG	ACA	ACG	ATC	GGA	GGA	CCG	AAG	GAG
136▶	Leu	Leu	Thr	Thr	Ile	Gly	Gly	Pro	Lys	Glu
4788	CTA	ACC	GCT	TTT	TTG	CAC	AAC	ATG	GGG	GAT
146▶	Leu	Thr	Ala	Phe	Leu	His	Asn	Met	Gly	Asp
4818	CAT	GTA	ACT	CGC	CTT	GAT	CGT	TGG	GAA	CCG
156▶	His	Val	Thr	Arg	Leu	Asp	Arg	Trp	Glu	Pro
4848	GAG	CTG	AAT	GAA	GCC	ATA	CCA	AAC	GAC	GAG
166▶	Glu	Leu	Asn	Glu	Ala	Ile	Pro	Asn	Asp	Glu
4878	CGT	GAC	ACC	ACG	ATG	CCT	GTA	GCA	ATG	GCA
176▶	Arg	Asp	Thr	Thr	Met	Pro	Val	Ala	Met	Ala
4908	ACA	ACG	TTG	CGC	AAA	CTA	TTA	ACT	GGC	GAA
186▶	Thr	Thr	Leu	Arg	Lys	Leu	Leu	Thr	Gly	Glu
4938	CTA	CTT	ACT	CTA	GCT	TCC	CGG	CAA	CAA	TTA
196▶	Leu	Leu	Thr	Leu	Ala	Ser	Arg	Gln	Gln	Leu

SUBSTITUTE SHEET (RULE 26)

63/65

Figure 14b (cont'd)

4968 ATA GAC TGG ATG GAG GCG GAT AAA GTT GCA  
 206▶ Ile Asp Trp Met Glu Ala Asp Lys Val Ala  
 4998 GGA CCA CTT CTG CGC TCG GCC CTT CCG GCT  
 216▶ Gly Pro Leu Leu Arg Ser Ala Leu Pro Ala  
 5028 GGC TGG TTT ATT GCT GAT AAA TCT GGA GCC  
 226▶ Gly Trp Phe Ile Ala Asp Lys Ser Gly Ala  
 5058 GGT GAG CGT GGG TCT CGC GGT ATC ATT GCA  
 236▶ Gly Glu Arg Gly Ser Arg Gly Ile Ile Ala  
 5088 GCA CTG GGG CCA GAT GGT AAG CCC TCC CGT  
 246▶ Ala Leu Gly Pro Asp Gly Lys Pro Ser Arg  
 5118 ATC GTA GTT ATC TAC ACG ACG GGG AGT CAG  
 256▶ Ile Val Val Ile Tyr Thr Thr Gly Ser Gln  
 5148 GCA ACT ATG GAT GAA CGA AAT AGA CAG ATC  
 266▶ Ala Thr Met Asp Glu Arg Asn Arg Gln Ile  
 5178 GCT GAG ATA GGT GCC TCA CTG ATT AAG CAT  
 276▶ Ala Glu Ile Gly Ala Ser Leu Ile Lys His  
 5208 TGG TAA CTGTCAGACCAAGTTTACTCATATATACTTT  
 286▶ Trp ...  
 5245 AGATTGATTTACGCGCCCTGTAGCGGCGCATTAAGCGCG  
 5284 GCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTT  
 5323 GCCAGCGCCCTAGCGCCCGCTCCTTTTCGCTTTTCTTCCCT  
 5362 TCCTTTTCTCGCCACGTTTCGCCGGCTTTCCCCGTCAAGCT  
 5401 CTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCT  
 5440 TTACGGCACCTCGACCCCAAAAAAAGTTGATTTGGGGTGAT  
 5479 GGTTCACGTAGTGGGCCATCGCCCTGATAGACGGTTT...

64/65

Figure 14b (cont'd)

5518 CGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGA  
5557 CTCTTGTTCCAAACTTGAACAACACTCAACCCTATCTCG  
5596 GGCTATTCTTTTGATTTATAAGGGATTTTGCCGATTTTCG  
5635 GCCTATTGGTTAAAAAATGAGCTGATTTAACAAAAATTT  
5674 AACGCGAATTTTAACAAAATATTAACGTTTACAATTTAA  
5713 AAGGATCTAGGTGAAGATCCTTTTTTGATAATCTCATGAC  
5752 CAAAATCCCTTAACGTGAGTTTTTCGTTCCACTGAGCGTC  
5791 AGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCC  
5830 TTTTTTTCTGCGCGTAATCTGCTGCTTGCAAACAAAAAA  
5869 ACCACCGCTACCAGCGGTGGTTTGTTTGCCGGATCAAGA  
5908 GCTACCAACTCTTTTTTCCGAAGGTAACTGGCTTCAGCAG  
5947 AGCGCAGATACCAAATACTGTCCTTCTAGTG TAGCCGTA  
5986 GTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTAC  
6025 ATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGC  
6064 CAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAG  
6103 ACGATAGTTACCGGATAAGGCGCAGCGGTCGGGCTGAAC  
6142 GGGGGGTTCGTGCACACAGCCCAGCTTGGAGCGAACGAC  
6181 CTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGA  
6220 AAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTA  
6259 TCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAG  
6298 GGAGCTTCCAGGGGGGAAACGCCTGGTATCTTTATAGTCC  
6337 TGTCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTT

SUBSTITUTE SHEET (RULE 26)

65/65

Figure 14b (cont'd)

6376 GTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAAACGC  
6415 CAGCAACGCGGCCTTTTTACGGTTCCTGGCCTTTTGCTG  
6454 GCCTTTTGCTCACATGTTCTTTCCTGCGTTATCCCCTGA  
6493 TTCTGTGGATAACCGTATTACCGCCTTTGAGTGAGCTGA  
6532 TACCGCTCGCCGCAGCCGAACGACCGAGCGCAGCGAGTC  
6571 AGTGAGCGAGGAAGCGGAAGAGCGCCTGATGCGGTATTT  
6610 TCTCCTTACGCATCTGTGCGGTATTTACACCCGCATAGG  
6649 GTCATGGCTGCGCCCCGACACCCGCCAACACCCGCTGAC  
6688 GCGCCCTGACGGGCTTGTCTGCTCCCGGCATCCGCTTAC  
6727 AGACAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTCAG  
6766 AGGTTTTTCACCGTCATCACCGAAACGCGCGAGGCAGCAA  
6805 GGAGATGGCGCCCAACAGTCCCCCGGCCACGGGGCCTGC  
6844 CACCATACCCACGCCGAAACAAGCGCTCATGAGCCCGAA  
6883 GTGGCGAGCCCGATCTTCCCCATCGGTGATGTCGGCGAT  
6922 ATAGGCGCCAGCAACCGCACCTGTGGCGCCGGTGATGCC  
6961 GGCCACGATGCGTCCGGCGTAGAGGATCTGCTCATGTTT  
7000 GACAGCTTATC

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(EMBL) (A) NAME: European Molecular Biology Laboratory  
(B) STREET: Meyerhofstrasse 1  
(C) CITY: Heidelberg  
(E) COUNTRY: DE  
(F) POSTAL CODE (ZIP): D-69117

(ii) TITLE OF INVENTION: Novel DNA Cloning Method

(iii) NUMBER OF SEQUENCES: 14

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30  
(EPO)

## (vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: EP 97121562.2  
(B) FILING DATE: 05-DEC-1997

## (vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: EP 98118756.0  
(B) FILING DATE: 05-OCT-1998

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6150 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: both  
(D) TOPOLOGY: both

## (vii) IMMEDIATE SOURCE:

(B) CLONE: pBAD24-recET

## (ix) FEATURE:

(A) NAME/KEY: misc\_feature  
(B) LOCATION: complement (96..974)  
(D) OTHER INFORMATION: /product= "araC"

## (ix) FEATURE:

(A) NAME/KEY: misc\_feature  
(B) LOCATION: 1320..2162  
(D) OTHER INFORMATION: /product= "t-recE"



## (ix) FEATURE:

- (A) NAME/KEY: misc\_feature
- (B) LOCATION: 2155..2972
- (D) OTHER INFORMATION: /product= "recT"

## (ix) FEATURE:

- (A) NAME/KEY: misc\_feature
- (B) LOCATION: 3493..4353
- (D) OTHER INFORMATION: /product= "bla"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```

ATCGATGCAT AATGTGCCTG TCAAATGGAC GAAGCAGGGA TTCTGCAAAC CCTATGCTAC      60
TCCGTCAAGC CGTCAATTGT CTGATTCGTT ACCAATTATG ACAACTTGAC GGCTACATCA      120
TTCACTTTTT CTTCACAACC GGCACGGAAC TCGCTCGGGC TGGCCCCGGT GCATTTTTTA      180
AATACCCGCG AGAAATAGAG TTGATCGTCA AAACCAACAT TGGACCGAC GGTGGCGATA      240
GGCATCCGGG TGGTGCTCAA AAGCAGCTTC GCCTGGCTGA TACGTTGGTC CTCGCGCCAG      300
CTTAAGACGC TAATCCCTAA CTGCTGGCGG AAAAGATGTG ACAGACGCGA CGGCGACAAG      360
CAAACATGCT GTGCGACGCT GGCATATCA AAATTGCTGT CTGCCAGGTG ATCGCTGATG      420
TACTGACAAG CCTCGCGTAC CCGATTATCC ATCGGTGGAT GGAGCGACTC GTTAATCGCT      480
TCCATGCGCC GCAGTAACAA TTGCTCAAGC AGATTTATCG CCAGCAGCTC CGAATAGCGC      540
CCTTCCCCTT GCCCGGCGTT AATGATTTC CCAAACAGGT CGCTGAAATG CGGCTGGTGC      600
GCTTCATCCG GSCGAAAGAA CCCCCTATTG GCAAATATTG ACGGCCAGTT AAGCCATTCA      660
TGCCAGTAGG CGCGCGGACG AAAGTAAACC CACTGGTGAT ACCATTGCGG AGCCTCCGGA      720
TGACGACCGT AGTGATGAAT CTCTCCTGGC GGAACAGCA AATATCACC CGGTGCGCAA      780
ACAAATTCTC GTCCCTGATT TTTACCACC CCCTGACCGC GAATGGTGAG ATTGAGAATA      840
TAACCTTTCA TTCCAGCGG TCGGTCGATA AAAAAATCGA GATAACCGTT GGCCTCAATC      900
GGCGTTAAAC CCGCCACCAG ATGGGCATTA AACGAGTATC CCGGCAGCAG GGGATCATTT      960
TGCGCTTCAG CCATACTTTT CATACTCCCG CCATTCAGAG AAGAAACCAA TTGTCCATAT     1020
TGCATCAGAC ATTGCCGTCA CTGCGTCTTT TACTGGCTCT TCTCGCTAAC CAAACCGGTA     1080
ACCCCGCTTA TTAAGCAT TCTGTAACAA AGCGGGACCA AAGCCATGAC AAAACGCGT     1140
AACAAAAGTG TCTATAATCA CGGCAGAAAA GTCCACATTG ATTATTTGCA CGGCGTCACA     1200
CTTTGCTATG CCATAGCATT TTTATCCATA AGATTAGCGG ATCCTACCTG ACGCTTTTTA     1260
TCGCAACTCT CTA CTGTTTCCATACCCG TTTTTTGGG CTAGCAGGAG GAATTCACCA     1320
TGGATCCCGT AATCGTAGAA GACATAGAGC CAGGTATTTA TTACGGAATT TCGAATGAGA     1380
ATTACCACGC GGSTCCCGGT ATCAGTAAGT CTCAGCTCGA TGACATTGCT GATACTCCGG     1440
CACTATATTT GTGGCGTAAA AATGCCCCCG TGGACACCAC AAAGACAAAA ACGCTCGATT     1500
TAGGAACTGC TTTCCACTGC CGGGTACTTG AACCAGGAAGA ATTCAGTAAC CGCTTTATCG     1560

```

TAGCACCTGA ATTTAACCGC CGTACAAACG CCGGAAAAGA AGAAGAGAAA GCGTTTCTGA	1620
TGGAATGCGC AAGCACAGGA AAAACGGTTA TCACTGCGGA AGAAGGCCGG AAAATTGAAC	1680
TCATGTATCA AAGCGTTATG GCTTTGCCGC TGGGGCAATG GCTTGTGAA AGCGCCGGAC	1740
ACGCTGAATC ATCAATTTAC TGGGAAGATC CTGAAACAGG AATTTTGTGT CCGTGCCGTC	1800
CGGACAAAAT TATCCCTGAA TTTCACTGGA TCATGGACGT GAAAACTACG GCGGATATTC	1860
AACGATTCAA AACCGCTTAT TACGACTACC GCTATCACGT TCAGGATGCA TTCTACAGTG	1920
ACGGTTATGA AGCACAGTTT GGAGTGCAGC CAACTTTCTG TTTTCTGGTT GCCAGCACAA	1980
CTATTGAATG CGGACGTTAT CCGGTTGAAA TTTTCATGAT GGGCGAAGAA GCAAACTGG	2040
CAGGTCAACA GGAATATCAC CGCAATCTGC GAACCCCTGTC TGAATGCCTG AATACCGATG	2100
AATGGCCAGC TATTAAGACA TTATCACTGC CCCGCTGGGC TAAGGAATAT GCAATGACT	2160
AAGCAACCAC CAATCGCAA AGCCGATCTG CAAAAACTC AGGGAAACCG TGCACCAGCA	2220
GCAGTTAAAA ATAGCGACGT GATTAGTTTT ATTAACCAGC CATCAATGAA AGAGCAACTG	2280
GCAGCAGCTC TTCCACGCCA TATGACGGCT GAACGTATGA TCCGTATCGC CACCACAGAA	2340
ATTGTAAG TTCCGGCGTT AGGAACTGT GACACTATGA GTTTTGTGAG TGCGATCGTA	2400
CAGTGTTCAC AGCTCGGACT TGAGCCAGGT AGCGCCCTCG GTCATGCATA TTTACTGCCT	2460
TTTGTAATA AAAACGAAAA GAGCGGTAAG AAGAACGTTT AGCTAATCAT TGGCTATCGC	2520
GGCATGATTG ATCTGGCTCG CCGTTCTGGT CAAATCGCCA GCCTGTCAGC CCGTGTGTC	2580
CGTGAAGGTG ACGAGTTTAG CTTCGAATTT GGCCTTGATG AAAAGTTAAT ACACCGCCCC	2640
GGAGAAAACG AAGATGCCCC GGTACCCAC GTCTATGCTG TCGCAAGACT GAAAGACGGA	2700
GGTACTCAGT TTGAAGTTAT GACGCGCAAA CAGATTGAGC TGGTGCGCAG CCTGAGTAAA	2760
GCTGGTAATA ACGGGCCGTG GGTAACCTAC TGGGAAGAAA TGGCAAAGAA AACGGCTATT	2820
CGTCGCCTGT TCAAATATTT GCGCGTATCA ATTGAGATCC AGCGTGCAGT ATCAATGGAT	2880
GAAAAGGAAC CACTGACAAT CGATCCTGCA GATTCCTCTG TATTAACCGG GGAATACAGT	2940
GTAATCGATA ATTCAGAGGA ATAGATCTAA GCTTGGCTGT TTTGGCGGAT GAGAGAAGAT	3000
TTTCAGCCTG ATACAGATTA AATCAGAACG CAGAAGCGGT CTGATAAAAC AGAATTTGCC	3060
TGGCGGCAGT AGCGCGGTGG TCCACCTGA CCCCATGCCG AACTCAGAAG TGAAACGCCG	3120
TAGCGCCGAT GGTAGTGTGG GGTCTCCCA TCGAGAGTA GGGAACTGCC AGGCATCAA	3180
TAAAACGAAA GGCTCAGTCG AAAGACTGGG CCTTTCTGTT TATCTGTTGT TTGTCGGTGA	3240
ACGCTCTCCT GAGTAGGACA AATCCGCCGG GAGCGGATTT GAACGTTGCG AAGCAACGGC	3300
CCGGAGGGTG GCGGGCAGGA CGCCCGCCAT AACTGCCAG GCATCAAATT AAGCAGAAGG	3360
CCATCCTGAC GGATGGCCTT TTGCGTTTC TACAACTCT TTTGTTTATT TTTCTAAATA	3420
CATTCAAATA TGTATCCGCT CATGAGACAA TAACCTGAT AAATGCTTCA ATAATATTGA	3480
AAAAGGAAGA GTATGAGTAT TCAACATTC CGTGTGCCC TTATTCCTT TTTTGGCGCA	3540
TTTTGCCTTC CTGTTTTTGC TCACCCAGAA ACGCTGGTGA AAGTAAAAGA TGCTGAAGAT	3600

CAGTTGGGTG CACGAGTGGG TTACATCGAA CTGGATCTCA ACAGCGGTAA GATCCTTGAG	3660
AGTTTTTCGCC CCGAAGAACG TTTTCCAATG ATGAGCACTT TTAAAGTTCT GCTATGTGGC	3720
GCGGTATTAT CCCGTGTTGA CGCCGGGCAA GAGCAACTCG GTCGCCGCAT ACACTATTCT	3780
CAGAATGACT TGGTTGAGTA CTCACCAGTC ACAGAAAAGC ATCTTACGGA TGGCATGACA	3840
GTAAGAGAAT TATGCAGTGC TGCCATAACC ATGAGTGATA ACACTGCGGC CAACTTACTT	3900
CTGACAACGA TCGGAGGACC GAAGGAGCTA ACCGCTTTTT TGCACAACAT GGGGGATCAT	3960
GTAAGTGGCC TTGATCGTTG GGAACCGGAG CTGAATGAAG CCATACCAA CGACGAGCGT	4020
GACACCACGA TGCCTGTAGC AATGGCAACA ACGTTGCGCA AACTATTAAC TGGCGAACTA	4080
CTTACTCTAG CTTCCCGGCA ACAATTAATA GACTGGATGG AGGCGGATAA AGTTGCAGGA	4140
CCACTTCTGC GCTCGGCCCT TCCGGCTGGC TGGTTTATTG CTGATAAATC TGGAGCCGGT	4200
GAGCGTGGGT CTCGCGGTAT CATTGCAGCA CTGGGGCCAG ATGGTAAGCC CTCCCGTATC	4260
GTAGTTATCT ACACGACGGG GAGTCAGGCA ACTATGGATG AACGAAATAG ACAGATCGCT	4320
GAGATAGGTG CCTCACTGAT TAAGCATTGG TAACTGTCAG ACCAAGTTTA CTCATATATA	4380
CTTTAGATTG ATTTACGCGC CCTGTAGCGG CGCATTAGC GCGGCGGGTG TGGTGGTTAC	4440
GCGCAGCGTG ACCGCTACAC TTGCCAGCGC CCTAGCGCCC GCTCCTTTCT CTTTCTTCCC	4500
TTCTTTTCTC GCCACGTTCT CCGGCTTTCC CCGTCAAGCT CTAAATCGGG GGCTCCCTTT	4560
AGGGTTCCGA TTTAGTGCTT TACGGCACCT CGACCCCAA AACTTGATT TGGGTGATGG	4620
TTACGTAAGT GGGCCATCGC CCTGATAGAC GGTTTTTTCT CCTTTGACGT TGGAGTCCAC	4680
GTCTTTTAAAT AGTGGACTCT TGTTCCAAAC TTGAACAACA CTCAACCCTA TCTCGGGCTA	4740
TTCTTTTGAT TTATAAGGGA TTTTGCCGAT TTCGGCCTAT TGGTTAAAAA ATGAGCTGAT	4800
TTAACAAAAA TTTAACGCGA ATTTTAACAA AATATTAAAG TTTACAATTT AAAAGGATCT	4860
AGGTGAAGAT CTTTTTTGAT AATCTCATGA CCAAATCCC TTAACGTGAG TTTTCGTTCC	4920
ACTGAGCGTC AGACCCCGTA GAAAAGATCA AAGGATCTTC TTGAGATCCT TTTTTCTGC	4980
GCGTAATCTG CTGCTTGCAA ACAAAAAAAC CACCGCTACC AGCGGTGGTT TGTGTGCCGG	5040
ATCAAGAGCT ACCAACTCTT TTTCCGAAGG TAACTGGCTT CAGCAGAGCG CAGATACCAA	5100
ATACTGTCTT TCTAGTGTAG CCGTAGTTAG GCCACCACTT CAAGAACTCT GTAGCACCGC	5160
CTACATACCT CGCTCTGCTA ATCCTGTTAC CAGTGGCTGC TGCCAGTGGC GATAAGTCGT	5220
GTCTTACCGG GTTGGACTCA AGACGATAGT TACCGGATAA GGCGCAGCGG TCGGGCTGAA	5280
CGGGGGGTTT GTGCACACAG CCCAGCTTGG AGCGAACGAC CTACACCGAA CTGAGATACC	5340
TACAGCGTGA GCTATGAGAA AGCGCCACGC TTCCCGAAGG GAGAAAGGCG GACAGGTATC	5400
CGGTAAGCGG CAGGGTCGGA ACAGGAGAGC GCACGAGGGA GCTTCCAGGG GGAAACGCTT	5460
GGTATCTTTA TAGTCCTGTC GGGTTTCGCC ACCTCTGACT TGAGCGTCGA TTTTGTGAT	5520
GCTCGTCAGG GGGGCGGAGC CTATGGAAA ACGCCAGCAA CGCGGCCTTT TTACGGTTCC	5580
TGGCCTTTTG CTGGCCTTTT GCTCACATGT TCTTCTCTGC GTTATCCCTT GATTCTGTGG	5640

ATAACCGTAT TACCGCCTTT GAGTGAGCTG ATACCGCTCG CCGCAGCCGA ACGACCGAGC 5700  
 GCAGCGAGTC AGTGAGCGAG GAAGCGGAAG AGCGCCTGAT GCGGTATTTT CTCCTTACGC 5760  
 ATCTGTGCGG TATTTACAC ACATAGGGT CATGGCTGCG CCCCACACC CGCCAACACC 5820  
 CGCTGACGGG CCTGACGGG CTTGTCTGCT CCGGCATCC GCTTACAGAC AAGCTGTGAC 5880  
 CGTCTCCGGG AGCTGCATGT GTCAGAGTT TTCACCGTCA TCACCGAAAC GCGCGAGGCA 5940  
 GCAAGGAGAT GCGGCCCAAC AGTCCCCCGG CCACGGGGCC TGCCACCATA CCCACGCCGA 6000  
 AACAAGCGCT CATGAGCCCG AAGTGGCGAG CCCGATCTTC CCCATCGGTG ATGTCGGCGA 6060  
 TATAGGCGCC AGCAACCGCA CCTGTGGCGC CGGTGATGCC GGCCACGATG CGTCCGGCGT 6120  
 AGAGGATCTG CTCATGTTTG ACAGCTTATC 6150

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 843 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(vii) IMMEDIATE SOURCE:

- (B) CLONE: t-recE

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..843
- (D) OTHER INFORMATION: /product= "t-recE"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

ATG GAT CCC GTA ATC GTA GAA GAC ATA GAG CCA GGT ATT TAT TAC GGA 48  
 Met Asp Pro Val Ile Val Glu Asp Ile Glu Pro Gly Ile Tyr Tyr Gly  
 1 5 10 15  
 ATT TCG AAT GAG AAT TAC CAC GCG GGT CCC GGT ATC AGT AAG TCT CAG 96  
 Ile Ser Asn Glu Asn Tyr His Ala Gly Pro Gly Ile Ser Lys Ser Gln  
 20 25 30  
 CTC GAT GAC ATT GCT GAT ACT CCG GCA CTA TAT TTG TGG CGT AAA AAT 144  
 Leu Asp Asp Ile Ala Asp Thr Pro Ala Leu Tyr Leu Trp Arg Lys Asn  
 35 40 45  
 GCC CCC GTG GAC ACC ACA AAG ACA AAA ACG CTC GAT TTA GGA ACT GCT 192  
 Ala Pro Val Asp Thr Thr Lys Thr Lys Thr Leu Asp Leu Gly Thr Ala  
 50 55 60  
 TTC CAC TGC CGG GTA CTT GAA CCG GAA GAA TTC AGT AAC CGC TTT ATC 240  
 Phe His Cys Arg Val Leu Glu Pro Glu Glu Phe Ser Asn Arg Phe Ile  
 65 70 75 80  
 GTA GCA CCT GAA TTT AAC CGC CGT ACA AAC GCC GGA AAA GAA GAA GAG 288  
 Val Ala Pro Glu Phe Asn Arg Arg Thr Asn Ala Gly Lys Glu Glu Glu  
 85 90 95  
 AAA GCG TTT CCG ATG GAA TGC GCA AGC ACA GGA AAA ACG GTT ATC ACT 336  
 Lys Ala Phe Leu Met Glu Cys Ala Ser Thr Gly Lys Thr Val Ile Thr  
 100 105 110

GCG GAA GAA GGC CGG AAA ATT GAA CTC ATG TAT CAA AGC GTT ATG GCT 384  
 Ala Glu Glu Gly Arg Lys Ile Glu Leu Met Tyr Gln Ser Val Met Ala  
 115 120 125  
 TTG CCG CTG GGG CAA TGG CTT GTT GAA AGC GCC GGA CAC GCT GAA TCA 432  
 Leu Pro Leu Gly Gln Trp Leu Val Glu Ser Ala Gly His Ala Glu Ser  
 130 135 140  
 TCA ATT TAC TGG GAA GAT CCT GAA ACA GGA ATT TTG TGT CGG TGC CGT 480  
 Ser Ile Tyr Trp Glu Asp Pro Glu Thr Gly Ile Leu Cys Arg Cys Arg  
 145 150 155 160  
 CCG GAC AAA ATT ATC CCT GAA TTT CAC TGG ATC ATG GAC GTG AAA ACT 528  
 Pro Asp Lys Ile Ile Pro Glu Phe His Trp Ile Met Asp Val Lys Thr  
 165 170 175  
 ACG GCG GAT ATT CAA CGA TTC AAA ACC GCT TAT TAC GAC TAC CGC TAT 576  
 Thr Ala Asp Ile Gln Arg Phe Lys Thr Ala Tyr Tyr Asp Tyr Arg Tyr  
 180 185 190  
 CAC GTT CAG GAT GCA TTC TAC AGT GAC GGT TAT GAA GCA CAG TTT GGA 624  
 His Val Gln Asp Ala Phe Tyr Ser Asp Gly Tyr Glu Ala Gln Phe Gly  
 195 200 205  
 GTG CAG CCA ACT TTC GTT TTT CTG GTT GCC AGC ACA ACT ATT GAA TGC 672  
 Val Gln Pro Thr Phe Val Phe Leu Val Ala Ser Thr Thr Ile Glu Cys  
 210 215 220  
 GGA CGT TAT CCG GTT GAA ATT TTC ATG ATG GGC GAA GAA GCA AAA CTG 720  
 Gly Arg Tyr Pro Val Glu Ile Phe Met Met Gly Glu Glu Ala Lys Leu  
 225 230 235 240  
 GCA GGT CAA CAG GAA TAT CAC CGC AAT CTG CGA ACC CTG TCT GAC TGC 768  
 Ala Gly Gln Gln Glu Tyr His Arg Asn Leu Arg Thr Leu Ser Asp Cys  
 245 250 255  
 CTG AAT ACC GAT GAA TGG CCA GCT ATT AAG ACA TTA TCA CTG CCC CGC 816  
 Leu Asn Thr Asp Glu Trp Pro Ala Ile Lys Thr Leu Ser Leu Pro Arg  
 260 265 270  
 TGG GCT AAG GAA TAT GCA AAT GAC TAA 843  
 Trp Ala Lys Glu Tyr Ala Asn Asp \*  
 275 280

## (2) INFORMATION FOR SEQ ID NO: 3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 281 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Met Asp Pro Val Ile Val Glu Asp Ile Glu Pro Gly Ile Tyr Tyr Gly  
 1 5 10 15  
 Ile Ser Asn Glu Asn Tyr His Ala Gly Pro Gly Ile Ser Lys Ser Gln  
 20 25 30  
 Leu Asp Asp Ile Ala Asp Thr Pro Ala Leu Tyr Leu Trp Arg Lys Asn  
 35 40 45  
 Ala Pro Val Asp Thr Thr Lys Thr Lys Thr Leu Asp Leu Gly Thr Ala  
 50 55 60

Phe His Cys Arg Val Leu Glu Pro Glu Glu Phe Ser Asn Arg Phe Ile  
 65 70 75 80  
 Val Ala Pro Glu Phe Asn Arg Arg Thr Asn Ala Gly Lys Glu Glu Glu  
 85 90 95  
 Lys Ala Phe Leu Met Glu Cys Ala Ser Thr Gly Lys Thr Val Ile Thr  
 100 105 110  
 Ala Glu Glu Gly Arg Lys Ile Glu Leu Met Tyr Gln Ser Val Met Ala  
 115 120 125  
 Leu Pro Leu Gly Gln Trp Leu Val Glu Ser Ala Gly His Ala Glu Ser  
 130 135 140  
 Ser Ile Tyr Trp Glu Asp Pro Glu Thr Gly Ile Leu Cys Arg Cys Arg  
 145 150 155 160  
 Pro Asp Lys Ile Ile Pro Glu Phe His Trp Ile Met Asp Val Lys Thr  
 165 170 175  
 Thr Ala Asp Ile Gln Arg Phe Lys Thr Ala Tyr Tyr Asp Tyr Arg Tyr  
 180 185 190  
 His Val Gln Asp Ala Phe Tyr Ser Asp Gly Tyr Glu Ala Gln Phe Gly  
 195 200 205  
 Val Gln Pro Thr Phe Val Phe Leu Val Ala Ser Thr Thr Ile Glu Cys  
 210 215 220  
 Gly Arg Tyr Pro Val Glu Ile Phe Met Met Gly Glu Glu Ala Lys Leu  
 225 230 235 240  
 Ala Gly Gln Gln Glu Tyr His Arg Asn Leu Arg Thr Leu Ser Asp Cys  
 245 250 255  
 Leu Asn Thr Asp Glu Trp Pro Ala Ile Lys Thr Leu Ser Leu Pro Arg  
 260 265 270  
 Trp Ala Lys Glu Tyr Ala Asn Asp \*  
 275 280

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 810 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(vii) IMMEDIATE SOURCE:

- (B) CLONE: rect

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..810
- (D) OTHER INFORMATION: /product= "rect"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

ATG ACT AAG CAA CCA CCA ATC GCA AAA ACC GAT CTG CAA AAA ACT CAG  
 Met Thr Lys Gln Pro Pro Ile Ala Lys Ala Asp Leu Gln Lys Thr Gln  
 285 290 295

GGA AAC CGT GCA CCA GCA GCA GTT AAA AAT AGC GAC GTG ATT AGT TTT Gly Asn Arg Ala Pro Ala Ala Val Lys Asn Ser Asp Val Ile Ser Phe 300 305 310	96
ATT AAC CAG CCA TCA ATG AAA GAG CAA CTG GCA GCA GCT CTT CCA CGC Ile Asn Gln Pro Ser Met Lys Glu Gln Leu Ala Ala Leu Pro Arg 315 320 325	144
CAT ATG ACG GCT GAA CGT ATG ATC CGT ATC GCC ACC ACA GAA ATT CGT His Met Thr Ala Glu Arg Met Ile Arg Ile Ala Thr Thr Glu Ile Arg 330 335 340 345	192
AAA GTT CCG GCG TTA GGA AAC TGT GAC ACT ATG AGT TTT GTC AGT GCG Lys Val Pro Ala Leu Gly Asn Cys Asp Thr Met Ser Phe Val Ser Ala 350 355 360	240
ATC GTA CAG TGT TCA CAG CTC GGA CTT GAG CCA GGT AGC GCC CTC GGT Ile Val Gln Cys Ser Gln Leu Gly Leu Glu Pro Gly Ser Ala Leu Gly 365 370 375	288
CAT GCA TAT TTA CTG CCT TTT GGT AAT AAA AAC GAA AAG AGC GGT AAA His Ala Tyr Leu Leu Pro Phe Gly Asn Lys Asn Glu Lys Ser Gly Lys 380 385 390	336
AAG AAC GTT CAG CTA ATC ATT GGC TAT CGC GGC ATG ATT GAT CTG GCT Lys Asn Val Gln Leu Ile Ile Gly Tyr Arg Gly Met Ile Asp Leu Ala 395 400 405	384
CGC CGT TCT GGT CAA ATC GCC AGC CTG TCA GCC CGT GTT GTC CGT GAA Arg Arg Ser Gly Gln Ile Ala Ser Leu Ser Ala Arg Val Val Arg Glu 410 415 420 425	432
GGT GAC GAG TTT AGC TTC GAA TTT GGC CTT GAT GAA AAG TTA ATA CAC Gly Asp Glu Phe Ser Phe Glu Phe Gly Leu Asp Glu Lys Leu Ile His 430 435 440	480
CGC CCG GGA GAA AAC GAA GAT GCC CCG GTT ACC CAC GTC TAT GCT GTC Arg Pro Gly Glu Asn Glu Asp Ala Pro Val Thr His Val Tyr Ala Val 445 450 455	528
GCA AGA CTG AAA GAC GGA GGT ACT CAG TTT GAA GTT ATG ACG CGC AAA Ala Arg Leu Lys Asp Gly Gly Thr Gln Phe Glu Val Met Thr Arg Lys 460 465 470	576
CAG ATT GAG CTG GTG CGC AGC CTG AGT AAA GCT GGT AAT AAC GGG CCG Gln Ile Glu Leu Val Arg Ser Leu Ser Lys Ala Gly Asn Asn Gly Pro 475 480 485	624
TGG GTA ACT CAC TGG GAA GAA ATG GCA AAG AAA ACG GCT ATT CGT CGC Trp Val Thr His Trp Glu Glu Met Ala Lys Lys Thr Ala Ile Arg Arg 490 495 500 505	672
CTG TTC AAA TAT TTG CCC GTA TCA ATT GAG ATC CAG CGT GCA GTA TCA Leu Phe Lys Tyr Leu Pro Val Ser Ile Glu Ile Gln Arg Ala Val Ser 510 515 520	720
ATG GAT GAA AAG GAA CCA CTG ACA ATC GAT CCT GCA GAT TCC TCT GTA Met Asp Glu Lys Glu Pro Leu Thr Ile Asp Pro Ala Asp Ser Ser Val 525 530 535	768
TTA ACC GGG GAA TAC AGT GTA ATC GAT AAT TCA GAG GAA TAG Leu Thr Gly Glu Tyr Ser Val Ile Asp Asn Ser Glu Glu * 540 545 550	810

## (2) INFORMATION FOR SEQ ID NO: 5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 270 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

```

Met Thr Lys Gln Pro Pro Ile Ala Lys Ala Asp Leu Gln Lys Thr Gln
 1           5           10           15
Gly Asn Arg Ala Pro Ala Ala Val Lys Asn Ser Asp Val Ile Ser Phe
          20           25           30
Ile Asn Gln Pro Ser Met Lys Glu Gln Leu Ala Ala Ala Leu Pro Arg
          35           40           45
His Met Thr Ala Glu Arg Met Ile Arg Ile Ala Thr Thr Glu Ile Arg
 50           55           60
Lys Val Pro Ala Leu Gly Asn Cys Asp Thr Met Ser Phe Val Ser Ala
 65           70           75           80
Ile Val Gln Cys Ser Gln Leu Gly Leu Glu Pro Gly Ser Ala Leu Gly
          85           90           95
His Ala Tyr Leu Leu Pro Phe Gly Asn Lys Asn Glu Lys Ser Gly Lys
          100          105          110
Lys Asn Val Gln Leu Ile Ile Gly Tyr Arg Gly Met Ile Asp Leu Ala
          115          120          125
Arg Arg Ser Gly Gln Ile Ala Ser Leu Ser Ala Arg Val Val Arg Glu
          130          135          140
Gly Asp Glu Phe Ser Phe Glu Phe Gly Leu Asp Glu Lys Leu Ile His
          145          150          155          160
Arg Pro Gly Glu Asn Glu Asp Ala Pro Val Thr His Val Tyr Ala Val
          165          170          175
Ala Arg Leu Lys Asp Gly Gly Thr Gln Phe Glu Val Met Thr Arg Lys
          180          185          190
Gln Ile Glu Leu Val Arg Ser Leu Ser Lys Ala Gly Asn Asn Gly Pro
          195          200          205
Trp Val Thr His Trp Glu Glu Met Ala Lys Lys Thr Ala Ile Arg Arg
          210          215          220
Leu Phe Lys Tyr Leu Pro Val Ser Ile Glu Ile Gln Arg Ala Val Ser
          225          230          235          240
Met Asp Glu Lys Glu Pro Leu Thr Ile Asp Pro Ala Asp Ser Ser Val
          245          250          255
Leu Thr Gly Glu Tyr Ser Val Ile Asp Asn Ser Glu Glu *
          260          265          270

```

## (2) INFORMATION FOR SEQ ID NO: 6:



## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 876 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: araC

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: complement (1..876)
- (D) OTHER INFORMATION: /product= "araC"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

```

TGACAACTTG ACGGCTACAT CATTCACTTT TTCTTCACAA CCGGCACGGA ACTCGCTCGG      60
GCTGGCCCCG GTGCATTTTT TAAATACCCG CGAGAAATAG AGTTGATCGT CAAAACCAAC      120
ATTGCGACCG ACGGTGGCGA TAGGCATCCG GGTGGTGCTC AAAAGCAGCT TCGCTGGCT      180
GATACGTTGG TCCTCGCGCC AGCTTAAGAC GCTAATCCCT AACTGCTGGC GGAAAAGATG      240
TGACAGACGC GACGGCGACA AGCAAACATG CTGTGCGACG CTGGCGATAT CAAAATTGCT      300
GTCTGCCAGG TGATCGCTGA TGTACTGACA AGCCTCGCGT ACCCGATTAT CCATCGGTGG      360
ATGGAGCGAC TCGTTAATCG CTTCCATGCG CCGCAGTAAC AATTGCTCAA GCAGATTTAT      420
CGCCAGCAGC TCCGAATAGC GCCCTTCCCC TTGCCCGGCG TTAATGATTT GCCCAAACAG      480
GTCGCTGAAA TCGGGCTGGT GCGCTTCATC CGGGCGAAAAG AACCCCGTAT TGGCAAATAT      540
TGACGGCCAG TTAAGCCATT CATGCCAGTA GGCGCGCGGA CGAAAGTAA CCGACTGGTG      600
ATACCATTGG CGAGCCTCCG GATGACGACC GTAGTGATGA ATCTCTCCTG GCGGGAACAG      660
CAAAATATCA CCGGTCGGC AAACAAATTC TCGTCCCTGA TTTTCACCA CCCCCTGACC      720
GCGAATGGTG AGATTGAGAA TATAACCTTT CATTCCCAGC GGTCCGTCGA TAAAAAAATC      780
GAGATAACCG TTGGCCTCAA TCGGCGTTAA ACCCGCCACC AGATGGGCAT TAAACGAGTA      840
TCCCGGCAGC AGGGGATCAT TTTGCGCTTC AGCCAT                                876

```

## (2) INFORMATION FOR SEQ ID NO: 7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 292 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

```

Met Ala Glu Ala Gln Asn Asp Pro Leu Leu Pro Gly Tyr Ser Phe Asn
 1           5           10           15
Ala His Leu Val Ala Gly Leu Thr Pro Ile Glu Ala Asn Gly Tyr Leu
          20           25           30

```

Asp Phe Phe Ile Asp Arg Pro Leu Gly Met Lys Gly Tyr Ile Leu Asn  
           35                          40                          45  
 Leu Thr Ile Arg Gly Gln Gly Val Val Lys Asn Gln Gly Arg Glu Phe  
           50                          55                          60  
 Val Cys Arg Pro Gly Asp Ile Leu Leu Phe Pro Pro Gly Glu Ile His  
           65                          70                          75                          80  
 His Tyr Gly Arg His Pro Glu Ala Arg Glu Trp Tyr His Gln Trp Val  
                           85                          90                          95  
 Tyr Phe Arg Pro Arg Ala Tyr Trp His Glu Trp Leu Asn Trp Pro Ser  
                           100                          105                          110  
 Ile Phe Ala Asn Thr Gly Phe Phe Arg Pro Asp Glu Ala His Gln Pro  
                           115                          120                          125  
 His Phe Ser Asp Leu Phe Gly Gln Ile Ile Asn Ala Gly Gln Gly Glu  
           130                          135                          140  
 Gly Arg Tyr Ser Glu Leu Leu Ala Ile Asn Leu Leu Glu Gln Leu Leu  
   145                          150                          155                          160  
 Leu Arg Arg Met Glu Ala Ile Asn Glu Ser Leu His Pro Pro Met Asp  
                           165                          170                          175  
 Asn Arg Val Arg Glu Ala Cys Gln Tyr Ile Ser Asp His Leu Ala Asp  
                           180                          185                          190  
 Ser Asn Phe Asp Ile Ala Ser Val Ala Gln His Val Cys Leu Ser Pro  
                           195                          200                          205  
 Ser Arg Leu Ser His Leu Phe Arg Gln Gln Leu Gly Ile Ser Val Leu  
   210                          215                          220  
 Ser Trp Arg Glu Asp Gln Arg Ile Ser Gln Ala Lys Leu Leu Leu Ser  
   225                          230                          235                          240  
 Thr Thr Arg Met Pro Ile Ala Thr Val Gly Arg Asn Val Gly Phe Asp  
                           245                          250                          255  
 Asp Gln Leu Tyr Phe Ser Arg Val Phe Lys Lys Cys Thr Gly Ala Ser  
                           260                          265                          270  
 Pro Ser Glu Phe Arg Ala Gly Cys Glu Glu Lys Val Asn Asp Val Ala  
                           275                          280                          285  
 Val Lys Leu Ser  
           290

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 861 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(vii) IMMEDIATE SOURCE:

- (B) CLONE: bla

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..861

(D) OTHER INFORMATION: /product= "bla"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

ATG AGT ATT CAA CAT TTC CGT GTC GCC CTT ATT CCC TTT TTT GCG GCA	48
Met Ser Ile Gln His Phe Arg Val Ala Leu Ile Pro Phe Phe Ala Ala	
295 300 305	
TTT TGC CTT CCT GTT TTT GCT CAC CCA GAA ACG CTG GTG AAA GTA AAA	96
Phe Cys Leu Pro Val Phe Ala His Pro Glu Thr Leu Val Lys Val Lys	
310 315 320	
GAT GCT GAA GAT CAG TTG GGT GCA CGA GTG GGT TAC ATC GAA CTG GAT	144
Asp Ala Glu Asp Gln Leu Gly Ala Arg Val Gly Tyr Ile Glu Leu Asp	
325 330 335 340	
CTC AAC AGC GGT AAG ATC CTT GAG AGT TTT CGC CCC GAA GAA CGT TTT	192
Leu Asn Ser Gly Lys Ile Leu Glu Ser Phe Arg Pro Glu Glu Arg Phe	
345 350 355	
CCA ATG ATG AGC ACT TTT AAA GTT CTG CTA TGT GGC GCG GTA TTA TCC	240
Pro Met Met Ser Thr Phe Lys Val Leu Leu Cys Gly Ala Val Leu Ser	
360 365 370	
CGT GTT GAC GCC GGG CAA GAG CAA CTC GGT CGC CGC ATA CAC TAT TCT	288
Arg Val Asp Ala Gly Gln Glu Leu Gly Arg Arg Ile His Tyr Ser	
375 380 385	
CAG AAT GAC TTG GTT GAG TAC TCA CCA GTC ACA GAA AAG CAT CTT ACG	336
Gln Asn Asp Leu Val Glu Tyr Ser Pro Val Thr Lys His Leu Thr	
390 395 400	
GAT GGC ATG ACA GTA AGA GAA TTA TGC AGT GCT GCC ATA ACC ATG AGT	384
Asp Gly Met Thr Val Arg Glu Leu Cys Ser Ala Ala Ile Thr Met Ser	
405 410 415 420	
GAT AAC ACT GCG GCC AAC TTA CTT CTG ACA ACG ATC GGA GGA CCG AAG	432
Asp Asn Thr Ala Ala Asn Leu Leu Leu Thr Thr Ile Gly Gly Pro Lys	
425 430 435	
GAG CTA ACC GCT TTT TTG CAC AAC ATG GGG GAT CAT GTA ACT CGC CTT	480
Glu Leu Thr Ala Phe Leu His Asn Met Gly Asp His Val Thr Arg Leu	
440 445 450	
GAT CGT TGG GAA CCG GAG CTG AAT GAA GCC ATA CCA AAC GAC GAG CGT	528
Asp Arg Trp Glu Pro Glu Leu Asn Glu Ala Ile Pro Asn Asp Glu Arg	
455 460 465	
GAC ACC ACG ATG CCT GTA GCA ATG GCA ACA ACG TTG CGC AAA CTA TTA	576
Asp Thr Thr Met Pro Val Ala Met Ala Thr Thr Leu Arg Lys Leu Leu	
470 475 480	
ACT GGC GAA CTA CTT ACT CTA GCT TCC CGG CAA CAA TTA ATA GAC TGG	624
Thr Gly Glu Leu Leu Thr Leu Ala Ser Arg Gln Gln Leu Ile Asp Trp	
485 490 495 500	
ATG GAG GCG GAT AAA GTT GCA GGA CCA CTT CTG CGC TCG GCC CTT CCG	672
Met Glu Ala Asp Lys Val Ala Gly Pro Leu Leu Arg Ser Ala Leu Pro	
505 510 515	
GCT GGC TGG TTT ATT GCT GAT AAA TCT GGA GCC GGT GAG CGT GGG TCT	720
Ala Gly Trp Phe Ile Ala Asp Lys Ser Gly Ala Gly Glu Arg Gly Ser	
520 525 530	

CGC GGT ATC ATT GCA GCA CTG GGG CCA GAT GGT AAG CCC TCC CGT ATC 768  
 Arg Gly Ile Ile Ala Ala Leu Gly Pro Asp Gly Lys Pro Ser Arg Ile  
 535 540 545

GTA GTT ATC TAC ACG ACG GGG AGT CAG GCA ACT ATG GAT GAA CGA AAT 816  
 Val Val Ile Tyr Thr Thr Gly Ser Gln Ala Thr Met Asp Glu Arg Asn  
 550 555 560

AGA CAG ATC GCT GAG ATA GGT GCC TCA CTG ATT AAG CAT TGG TAA 861  
 Arg Gln Ile Ala Glu Ile Gly Ala Ser Leu Ile Lys His Trp \*  
 565 570 575

## (2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 287 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Met Ser Ile Gln His Phe Arg Val Ala Leu Ile Pro Phe Phe Ala Ala  
 1 5 10 15

Phe Cys Leu Pro Val Phe Ala His Pro Glu Thr Leu Val Lys Val Lys  
 20 25 30

Asp Ala Glu Asp Gln Leu Gly Ala Arg Val Gly Tyr Ile Glu Leu Asp  
 35 40 45

Leu Asn Ser Gly Lys Ile Leu Glu Ser Phe Arg Pro Glu Glu Arg Phe  
 50 55 60

Pro Met Met Ser Thr Phe Lys Val Leu Leu Cys Gly Ala Val Leu Ser  
 65 70 75 80

Arg Val Asp Ala Gly Gln Glu Gln Leu Gly Arg Arg Ile His Tyr Ser  
 85 90 95

Gln Asn Asp Leu Val Glu Tyr Ser Pro Val Thr Glu Lys His Leu Thr  
 100 105 110

Asp Gly Met Thr Val Arg Glu Leu Cys Ser Ala Ala Ile Thr Met Ser  
 115 120 125

Asp Asn Thr Ala Ala Asn Leu Leu Thr Thr Ile Gly Gly Pro Lys  
 130 135 140

Glu Leu Thr Ala Phe Leu His Asn Met Gly Asp His Val Thr Arg Leu  
 145 150 155 160

Asp Arg Trp Glu Pro Glu Leu Asn Glu Ala Ile Pro Asn Asp Glu Arg  
 165 170 175

Asp Thr Thr Met Pro Val Ala Met Ala Thr Thr Leu Arg Lys Leu Leu  
 180 185 190

Thr Gly Glu Leu Leu Thr Leu Ala Ser Arg Gln Gln Leu Ile Asp Trp  
 195 200 205

Met Glu Ala Asp Lys Val Ala Gly Pro Leu Leu Arg Ser Ala Leu Pro  
 210 215 220

Ala Gly Trp Phe Ile Ala Asp Lys Ser Gly Ala Gly Glu Arg Gly Ser

225		230		235		240
Arg Gly Ile Ile	Ala Ala Leu Gly Pro	Asp Gly Lys Pro Ser	Arg Ile			
	245	250	255			
Val Val Ile Tyr Thr Thr Gly Ser	Gln Ala Thr Met Asp Glu Arg Asn					
	260	265	270			
Arg Gln Ile Ala Glu Ile Gly Ala Ser Leu Ile Lys His Trp						
	275	280	285			

## (2) INFORMATION FOR SEQ ID NO: 10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7195 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: pBAD-ETgamma

## (ix) FEATURE:

- (A) NAME/KEY: misc\_feature
- (B) LOCATION: 3588..74004
- (D) OTHER INFORMATION: /product= "red gamma"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

ATCGATGCAT AATGTGCCTG TCAAATGGAC GAAGCAGGGA TTCTGCAAAC CCTATGCTAC	60
TCCGTCAAGC CGTCAATTGT CTGATTCGTT ACCAATTATG ACAACTTGAC GGCTACATCA	120
TTCACTTTTT CTTCACAACC GGCACGGAAC TCGCTCGGGC TGGCCCCGGT GCATTTTTTTA	180
AATACCCGCG AGAAATAGAG TTGATCGTCA AAACCAACAT TGCGACCGAC GGTGGCGATA	240
GGCATCCGGG TGGTGCTCAA AAGCAGCTTC GCCTGGCTGA TACGTTGGTC CTCGCGCCAG	300
CTTAAGACGC TAATCCCTAA CTGCTGGCGG AAAAGATGTG ACAGACGCGA CGGCGACAAG	360
CATACATGCT GTGCGACGCT GCGGATATCA AATTGCTGT CTGCCAGGTG ATCGCTGATG	420
TACTGACAAG CCTCGCGTAC CCGATTATCC ATCGGTGGAT GGAGCGACTC GTTAATCGCT	480
TCCATGCGCC GCAGTAACAA TTGCTCAAGC AGATTTATCG CCAGCAGCTC CGAATAGCGC	540
CCTTCCCTTT GCCCGGCGTT AATGATTTCG CCAAACAGGT CGCTGAAATG CGGCTGGTGC	600
GCTTCATCCG GCGGAAAGAA CCCCCTATTG GCAAATATTG ACGGCCAGTT AAGCCATTCA	660
TGCCAGTAGG CGCGCGGACG AAAGTAAACC CACTGGTGAT ACCATTCGCG AGCCTCCGGA	720
TGACGACCGT AGTGATGAAT CTCTCCTGGC GGGAACAGCA AAATATCACC CGGTCGGCAA	780
ACAAATTCTC GTCCCTGATT TTTCACCACC CCTGACCGC GAATGGTGAG ATTGAGAATA	840
TAACCTTTCA TTCCCAGCGG TCGGTCGATA AAAAAATCGA GATAACCGTT GGCCTCAATC	900
GGCGTTAAAC CCGCCACCAG ATGGGCATTA AACGAGTATC CCGGCAGCAG GGGATCATTT	960
TGCGCTTCAG CCATACTTTT CATACTCCCG CCATTCAGAG AAGAAACCAA TTGTCCATAT	1020

TGCATCAGAC ATTGCCGTCA CTGCGTCTTT TACTGGCTCT TCTCGCTAAC CAAACCGGTA	1080
ACCCCGCTTA TTAAGCAT TCTGTAACAA AGCGGGACCA AAGCCATGAC AAAAACGCGT	1140
AACAAAAGTG TCTATAATCA CGGCAGAAAA GTCCACATTG ATTATTTGCA CGGCGTCACA	1200
CTTTGCTATG CCATAGCATT TTTATCCATA AGATTAGCGG ATCCTACCTG ACGCTTTTTA	1260
TCGCAACTCT CTACTGTTTC TCCATACCCG TTTTTTTGGG CTAGCAGGAG GAATTCACCA	1320
TGGATCCCGT AATCGTAGAA GACATAGAGC CAGGTATTTA TTACGGAATT TCGAATGAGA	1380
ATTACCACGC GGGTCCCGGT ATCAGTAAGT CTCAGCTCGA TGACATTGCT GATACTCCGG	1440
CACTATAATT GTGGGTAAA AATGCCCCCG TGGACACCAC AAAGACAAAA ACGCTCGATT	1500
TAGGAACTGC TTTCCACTGC CGGCTACTTG AACCGGAAGA ATTCAGTAAC CGCTTTATCG	1560
TAGCACCTGA ATTTAACCGC CGTACAAACG CCGGAAAAGA AGAAGAGAAA GCGTTTCTGA	1620
TGGAATGCGC AAGCACAGGA AAAACGGTTA TCACTGCGGA AGAAGGCCGG AAAATTGAAC	1680
TCATGTATCA AAGCGTTATG GCTTTGCCGC TGGGGCAATG GCTTGTGAA AGCGCCGGAC	1740
ACGCTGAATC ATCAATTTAC TGGGAAGATC CTGAAACAGG AATTTTGTGT CGGTGCCGTC	1800
CGGACAAAAT TATCCCTGAA TTTCACTGGA TCATGGACGT GAAACTACG GCGGATATTC	1860
AACGATTCAA AACCGCTTAT TACGACTACC GCTATCACGT TCAGGATGCA TTCTACAGTG	1920
ACGGTTATGA AGCACAGTTT GGAGTGCAGC CAACTTTCTG TTTTCTGGTT GCCAGCACAA	1980
CTATTGAATG CGGACGTTAT CCGGTTGAAA TTTTCATGAT GGGCGAAGAA GCAAACTGG	2040
CAGGTCAACA GGAATATCAC CGCAATCTGC GAACCCTGTC TGACTGCCTG AATACCGATG	2100
AATGCCCAGC TATTAAGACA TTATCACTGC CCCGCTGGGC TAAGGAATAT GCAAATGACT	2160
AGATCTCGAG GTACCCGAGC ACGTGTGAC AATTAATCAT CGGCATAGTA TATCGGCATA	2220
GTATAATACG ACAAGGTGAG GAACTAAACC ATGGCTAAGC AACCACCAAT CGCAAAAGCC	2280
GATCTGCAAA AACTCAGGG AAACCGTGCA CCAGCAGCAG TTAAAAATAG CGACGTGATT	2340
AGTTTTATTA ACCAGCCATC AATGAAAGAG CAACTGGCAG CAGCTCTTCC ACGCCATATG	2400
ACGGCTGAAC GTATGATCCG TATCGCCACC ACAGAAATTC GTAAAGTTCC GCGGTTAGGA	2460
AACTGTGACA CTATGAGTTT TGTCAGTGCG ATCGTACAGT GTTCACAGCT CGGACTTGAG	2520
CCAGGTAGCG CCCTCGGTCA TGCATATTTA CTGCCTTTTG GTAATAAAAA CGAAAAGAGC	2580
GGTAAAAAGA ACGTTCAGCT AATCATTGGC TATCGCGGCA TGATTGATCT GGCTCGCCGT	2640
TCTGGTCAAA TCGCCAGCCT GTCAGCCCGT GTTGTCCGTG AAGGTGACGA GTTTAGCTTC	2700
GAATTTGGCC TTGATGAAAA GTTAATACAC CGCCCGGGAG AAAACGAAGA TGCCCCGGTT	2760
ACCCACGTCT ATGCTGTGCG AAGACTGAAA GACGGAGGTA CTCAGTTTGA AGTTATGACG	2820
CGCAAACAGA TTGAGCTGGT GCGCAGCCTG AGTAAAGCTG GTAATAACGG GCCGTGGGTA	2880
ACTCACTGGG AAGAAATGGC AAAGAAAACG GCTATTCGTC GCCTGTTCAA ATATTTGCCC	2940
GATCAATTG AGATCCAGCG TGCAGTATCA ATGGATGAAA AGGAACCACT GACAATCGAT	3000
CCTGCAGATT CCTCTGTATT AACCGGGGAA TACAGTGTA TCGATAATTC AGAGGAATAG	3060

ATCTAAGCTT CCTGCTGAAC ATCAAAGGCA AGAAAACATC TGTTGTCAAA GACAGCATCC	3120
TTGAACAAGG ACAATTAACA GTTAACAAAT AAAAACGCAA AAGAAAATGC CGATATCCTA	3180
TTGGCATTTC CTTTTATTTC TTATCAACAT AAAGGTGAAT CCCATACCTC GAGCTTCACG	3240
CTGCCGCAAG CACTCAGGGC GCAAGGGCTG CTAAAAGGAA GCGGAACACG TAGAAAGCCA	3300
GTCCGCAGAA ACGGTGCTGA CCCCAGATGA ATGTCAGCTA CTGGGCTATC TGGACAAGGG	3360
AAAACGCAAG CGCAAAGAGA AAGCAGGTAG CTTCAGTGG GCTTACATGG CGATAGCTAG	3420
ACTGGGCGGT TTTATGGACA GCAAGCGAAC CGGAATTGCC AGCTGGGGCG CCCTCTGGTA	3480
AGGTTGGGAA GCCCTGCAA GTAAACTGGA TGGCTTTCTT GCCGCCAAGG ATCTGATGGC	3540
GCAGGGGATC AAGATCTGAT CAAGAGACAG GATGAGGATC GTTTCGCATG GATATTAATA	3600
CTGAAACTGA GATCAAGCAA AAGCATTAC TAACCCCTT TCCTGTTTTT CTAATCAGCC	3660
CGGCATTTTC CGGGCGATAT TTTACAGCT ATTTCAGGAG TTCAGCCATG AACGCTTATT	3720
ACATTCAGGA TCGTCTTGAG GCTCAGAGCT GGGCGCGTCA CTACCAGCAG CTCGCCCCGTG	3780
AAGAGAAAGA GGCAGAACTG GCAGACGACA TGGAAAAAGG CCTGCCCCAG CACCTGTTTG	3840
AATCGCTATG CATCGATCAT TTGCAACGCC ACGGGGCCAG CAAAAAATCC ATTACCCGTG	3900
CGTTTGATGA CGATGTTGAG TTTCAGGAGC GCATGGCAGA ACACATCCGG TACATGGTTG	3960
AAACCATTCG TCACCACCAG GTTGATATTG ATTCAGAGGT ATAAAACGAG TAGAAGCTTG	4020
GCTGTTTTTG CGGATGAGAG AAGATTTTCA GCCTGATACA GATTAAATCA GAACGCAGAA	4080
GCGGTCTGAT AAAACAGAAT TTGCCTGGCG GCAGTAGCGC GGTGGTCCCA CCTGACCCCA	4140
TGCCGAATC AGAAGTGAAG CGCCGTAGCG CCGATGGTAG TGTGGGGTCT CCCCATGCGA	4200
GAGTAGGGAA CTGCCAGGCA TCAAATAAAA CGAAGGCTC AGTCGAAAGA CTGGGCCTTT	4260
CGTTTTATCT GTTGTGTTGTC GGTGAACGCT CTCCTGAGTA GGACAAATCC GCCGGGAGCG	4320
GATTTGAACG TTGCGAAGCA ACGGCCCGGA GGGTGGCGGG CAGGACGCCC GCCATAAACT	4380
GCCAGGCATC AAATTAAGCA GAAGGCCATC CTGACGGATG GCCTTTTTTG GTTTCTACAA	4440
ACTCTTTTGT TTATTTTTCT AAATACATTC AAATATGTAT CCGCTCATGA GACAATAACC	4500
CTGATAAATG CTTCAATAAT ATTGAAAAAG GAAGAGTATG AGTATTCAAC ATTTCCGTGT	4560
CGCCCTTATT CCCTTTTTTG CGGCATTTTG CCTTCCTGTT TTTGCTCACC CAGAAACGCT	4620
GGTGAAAGTA AAAGATGCTG AAGATCAGTT GGGTGCACGA GTGGGTACCA TCGAACTGGA	4680
TCTCAACAGC GGTAAAGATCC TTGAGAGTTT TCGCCCCGAA GAACGTTTTT CAATGATGAG	4740
CACTTTTAAA GTTCTGCTAT GTGGCGCGGT ATTATCCCGT GTTGACGCCG GGCAAGAGCA	4800
ACTCGGTCGC CGCATACACT ATTCTCAGAA TGACTTGTTT GAGTACTCAC CAGTCACAGA	4860
AAAGCATCTT ACGGATGGCA TGACAGTAAG AGAATTATGC AGTGCTGCCA TAACCATGAG	4920
TGATAACACT GCGGCCAACT TACTTCTGAG AATGATGGA GGACCGAAGG AGCTAACCGC	4980
TTTTTTGCAC AACATGGGGG ATCATGTAAT TCCCTTGAT CGTTGGGAAC CGGAGCTGAA	5040
TGAAGCCATA CCAAACGACG AGCGTGACAC CAGGATGCCT GTAGCAATGG CAACAACGTT	5100

GCGCAAACTA TTAAGTGGCG AACTACTTAC TCTAGCTTCC CGGCAACAAT TAATAGACTG	5160
GATGGAGGCG GATAAAGTTG CAGGACCACT TCTGCGCTCG GCCCTTCCGG CTGGCTGGTT	5220
TATTGCTGAT AAATCTGGAG CCGGTGAGCG TGGGTCTCGC GGTATCATTG CAGCACTGGG	5280
GCCAGATGGT AAGCCCTCCC GTATCGTAGT TATCTACACG ACGGGGAGTC AGGCAACTAT	5340
GGATGAACGA AATAGACAGA TCGCTGAGAT AGGTGCCTCA CTGATTAAGC ATTGGTAACT	5400
GTCAGACCAA GTTTACTCAT ATATACTTTA GATTGATTTA CGCGCCCTGT AGCGGCGCAT	5460
TAAGCGCGGC GGGTGTGGTG GTTACGCGCA GCGTGACCGC TACACTTGCC AGC3CCCTAG	5520
CGCCCGCTCC ITTCGCTTTC TTCCCTTCCT TTCTCGCCAC GTTCGCCGGC TTTCCCGTC	5580
AAGCTCTAAA TCGGGGGCTC CCTTTAGGGT TCCGATTTAG TGCTTTACGG CACCTCGACC	5640
CCAAAAAAGT TGATTTGGGT GATGGTTCAC GTAGTGGGCC ATCGCCCTGA TAGACGGTTT	5700
TTCCGCCCTT GACGTTGGAG TCCACGTTCT TTAATAGTGG ACTCTTGTTT CAACTTGAA	5760
CAACACTCAA CCTATCTCG GGCTATTCTT TTGATTTATA AGGGATTTTG CCGATTTCCG	5820
CCTATTGGTT AAAAAATGAG CTGATTTAAC AAAAATTTAA CGCGAATTTT AACAAAATAT	5880
TAACGTTTAC AATTTAAAAG GATCTAGGTG AAGATCCTTT TTGATAATCT CATGACCAAA	5940
ATCCCTTAAC GTGAGTTTTT GTTCCACTGA GCGTCAGACC CCGTAGAAAA GATCAAAGGA	6000
TCTTCTTGAG ATCCTTTTTT TCTGCGCGTA ATCTGCTGCT TGCAAACAAA AAAACCACCG	6060
CTACCAGCGG TGGTTTGTTC GCCGGATCAA GAGCTACCAA CTCTTTTTTC GAAGGTAAGT	6120
GGCTTCAGCA GAGCGCAGAT ACCAAATACT GTCCTTCTAG TGTAGCCGTA GTTAGGCCAC	6180
CACTTCAAGA ACTCTGTAGC ACCGCCTACA TACCTCGCTC TGCTAATCCT GTTACCAGTG	6240
GCTGCTGCCA GTGGCGATAA GTCGTGCTT ACCGGGTGG ACTCAAGACG ATAGTTACCG	6300
GATAAGGCGC AGCGGTGCGG CTGAACGGGG GGTTCGTGCA CACAGCCCAG CTTGGAGCGA	6360
ACGACCTACA CGGAAGTGA ATACCTACAG CGTGAGCTAT GAGAAAGCGC CACGCTTCCC	6420
GAAGGGAGAA AGGCGGACAG GTATCCGGTA AGCGGCAGGG TCGGAACAGG AGAGCGCAGC	6480
AGGGAGCTTC CAGGGGGAAA CGCCTGGTAT CTTTATAGTC CTGTGCGGTT TCGCCACCTC	6540
TGACTTGAGC GTCGATTTTT GTGATGCTCG TCAGGGGGGC GGAGCCTATG GAAAAACGCC	6600
AGCAACGCGG CCTTTTTACG GTTCTGGCC TTTTGCTGGC CTTTGCTCA CATGTTCTTT	6660
CCTGCGTTAT CCCCTGATTC TGTGGATAAC CGTATTACCG CCTTTGAGTG AGCTGATACC	6720
GCTCGCCGCA GCCGAACGAC CGAGCGCAGC GAGTCAGTGA GCGAGGAAGC GGAAGAGCGC	6780
CTGATGCGGT ATTTTCTCCT TACGCATCTG TCGGGTATTT CACACCGCAT AGGGTCATGG	6840
CTGCGCCCCG ACACCCGCCA ACACCCGCTG ACGCGCCCTG ACGGGCTTGT CTGCTCCCGG	6900
CATCCGCTTA CAGACAAGCT GTGACCGTCT CCGGGAGCTG CATGTGTCAG AGGTTTTTAC	6960
CGTCATCACC GAAACGCGCG AGGCAGCAAG GAGATGGGCG CCAACAGTCC CCCGGCCACG	7020
GGGCCTGCCA CCATACCCAC GCCGAAACAA GCGCTCATGA GCCCGAAGTG GCGAGCCCGA	7080
TCTTCCCCAT CGGTGATGTC GGCGATATAG GCGCCAGCAA CCGCACCTGT GGCGCCGGTG	7140



ATGCCGGCCA CGATGCGTCC GCGGTAGAGG ATCTGCTCAT GTTTGACAGC TTATC

7195

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7010 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(vii) IMMEDIATE SOURCE:

- (B) CLONE: pBAD-alpha-beta-gamma

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1320..2000
- (D) OTHER INFORMATION: /product= "red alpha"

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 2086..2871
- (D) OTHER INFORMATION: /product= "red beta"

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 3403..3819
- (D) OTHER INFORMATION: /product= "red gamma"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

ATCGATGCAT AATGTGCTG TCAAATGGAC GAAGCAGGGA TTCTGCAAAC CCTATGCTAC	60
TCCGTCAAGC CGTCATTGT CTGATTCGTT ACCAATTATG ACAACTTGAC GGCTACATCA	120
TTCACTTTTTT CTTCAACAACC GGCACGGAAC TCGCTCGGGC TGGCCCCGGT GCATTTTTTA	180
AATACCCGCG AGAATAGAG TTGATCGTCA AAACCAACAT TCGACCGAC GGTGGCGATA	240
GGCATCCGGG TGGTGCTCAA AAGCAGCTTC GCCTGGCTGA TACGTTGGTC CTCGCGCCAG	300
CTTAAGACGC TAATCCCTAA CTGCTGGCGG AAAAGATGTG ACAGACGCGA CGGCGACAAG	360
CAAACATGCT GTGCGACGCT GCGGATATCA AAATTGCTGT CTGCCAGGTG ATCGCTGATG	420
TACTGACAAG CCTCGCGTAC CCGATTATCC ATCGGTGGAT GGAGCGACTC GTTAATCGCT	480
TCCATGCGCC GCAGTAACAA TTGCTCAAGC AGATTATCG CCAGCAGCTC CGAATAGCGC	540
CCTTCCCCTT GCCCGGCGTT AATGATTTGC CCAAACAGGT CGCTGAAATG CGGCTGGTGC	600
GCTTCATCCG GCGGAAAGAA CCGGTATTG GCAAATATTG ACGGCCAGTT AAGCCATTCA	660
TGCCAGTAGG CGCGCGGACG AAAGTAAACC CACTGGTGAT ACCATTGCGG AGCCTCCGGA	720
TGACGACCGT AGTGATGAAT CTCTGCTGGC GGGAACAGCA AAATATCACC CGGTGCGCAA	780
ACAAATTCTC GTCCCTGATT TTTCACCACC CCCTGACCGC GAATGGTGAG ATTGAGAATA	840
TAACCTTTCA TTCCGAGCGG TCGGTCGATA AAAAAATCGA GATAACCGTT GGCCTCAATC	900
GCGGTAAAC CCGCCACCAG ATGGGCATTA AACGAGTATC CCGGCAGCAG GGGATCATTT	960

TGCGCTTCAG CCATACTTTT CATACTCCCG CCATTTCAGAG AAGAAACCAA TTGTCCATAT	1020
TGCATCAGAC ATTGCCGTCA CTGCGTCTTT TACTGGCTCT TCTCGCTAAC CAAACCGGTA	1080
ACCCCGCTTA TTAAGCAT TCTGTAACAA AGCGGGACCA AAGCCATGAC AAAACGCGT	1140
AACAAAAGTG TCTATAATCA CGGCAGAAAA GTCCACATTG ATTATTTGCA CGGCGTCACA	1200
CTTTGCTATG CCATAGCATT TTTATCCATA AGATTAGCGG ATCCTACCTG ACGCTTTTTA	1260
TCGCAACTCT CTACTGTTTC TCCATACCCG TTTTTTTGGG CTAGCAGGAG GAATTCACC	1319
ATG ACA CCG GAC ATT ATC CTG CAG CGT ACC GGG ATC GAT GTG AGA GCT Met Thr Pro Asp Ile Ile Leu Gln Arg Thr Gly Ile Asp Val Arg Ala 290 295 300	1367
GTC GAA CAG GGG GAT GAT GCG TGG CAC AAA TTA CGG CTC GGC GTC ATC Val Glu Gln Gly Asp Asp Ala Trp His Lys Leu Arg Leu Gly Val Ile 305 310 315	1415
ACC GCT TCA GAA GTT CAC AAC GTG ATA GCA AAA CCC CGC TCC GGA AAG Thr Ala Ser Glu Val His Asn Val Ile Ala Lys Pro Arg Ser Gly Lys 320 325 330 335	1463
AAG TGG CCT GAC ATG AAA ATG TCC TAC TTC CAC ACC CTG CTT GCT GAG Lys Trp Pro Asp Met Lys Met Ser Tyr Phe His Thr Leu Leu Ala Glu 340 345 350	1511
GTT TGC ACC GGT GTG GCT CCG GAA GTT AAC GCT AAA GCA CTG GCC TGG Val Cys Thr Gly Val Ala Pro Glu Val Asn Ala Lys Ala Leu Ala Trp 355 360 365	1559
GGA AAA CAG TAC GAG AAC GAC GCC AGA ACC CTG TTT GAA TTC ACT TCC Gly Lys Gln Tyr Glu Asn Asp Ala Arg Thr Leu Phe Glu Phe Thr Ser 370 375 380	1607
GGC GTG AAT GTT ACT GAA TCC CCG ATC ATC TAT CGC GAC GAA AGT ATG Gly Val Asn Val Thr Glu Ser Pro Ile Ile Tyr Arg Asp Glu Ser Met 385 390 395	1655
CGT ACC GCC TGC TCT CCC GAT GGT TTA TGC AGT GAC GGC AAC GGC CTT Arg Thr Ala Cys Ser Pro Asp Gly Leu Cys Ser Asp Gly Asn Gly Leu 400 405 410 415	1703
GAA CTG AAA TGC CCG TTT ACC TCC CGG GAT TTC ATG AAG TTC CGG CTC Glu Leu Lys Cys Pro Phe Thr Ser Arg Asp Phe Met Lys Phe Arg Leu 420 425 430	1751
GGT GGT TTC GAG GCC ATA AAG TCA GCT TAC ATG GCC CAG GTG CAG TAC Gly Gly Phe Glu Ala Ile Lys Ser Ala Tyr Met Ala Gln Val Gln Tyr 435 440 445	1799
AGC ATG TGG GTG ACG CGA AAA AAT GCC TGG TAC TTT GCC AAC TAT GAC Ser Met Trp Val Thr Arg Lys Asn Ala Trp Tyr Phe Ala Asn Tyr Asp 450 455 460	1847
CCG CGT ATG AAG CGT GAA GGC CTG CAT TAT GTC GTG ATT GAG CGG GAT Pro Arg Met Lys Arg Glu Gly Leu His Tyr Val Ile Glu Arg Asp 465 470 475	1895
GAA AAG TAC ATG GCG AGT TTT GAC GAG ATC GTG CCG GAG TTC ATC GAA Glu Lys Tyr Met Ala Ser Phe Asp Glu Ile Val Pro Glu Phe Ile Glu 480 485 490 495	1943
AAA ATG GAC GAG GCA CTG GCT GAA ATT GGT TTT GTA TTT GGG GAG CAA Lys Met Asp Glu Ala Leu Ala Glu Ile Gly Phe Val Phe Gly Glu Gln 500 505 510	1991

TGG CGA TAG ATCCGGTACC CGAGCACGTG TTGACAATTA ATCATCGGCA	2040
Trp Arg *	
TAGTATATCG GCATAGTATA ATACGACAAG GTGAGGAACT AAACC ATG AGT ACT	2094
Met Ser Thr	
1	
GCA CTC GCA ACG CTG GCT GGG AAG CTG GCT GAA CGT GTC GGC ATG GAT	2142
Ala Leu Ala Thr Leu Ala Gly Lys Leu Ala Glu Arg Val Gly Met Asp	
5 10 15	
TCT GTC GAC CCA CAG GAA CTG ATC ACC ACT CTT CGC CAG ACG GCA TTT	2190
Ser Val Asp Pro Gln Glu Leu Ile Thr Thr Leu Arg Gln Thr Ala Phe	
20 25 30 35	
AAA GGT GAT GCC AGC GAT GCG CAG TTC ATC GCA TTA CTG ATC GTT GCC	2238
Lys Gly Asp Ala Ser Asp Ala Gln Phe Ile Ala Leu Leu Ile Val Ala	
40 45 50	
AAC CAG TAC GGC CTT AAT CCG TGG ACG AAA GAA ATT TAC GCC TTT CCT	2286
Asn Gln Tyr Gly Leu Asn Pro Trp Thr Lys Glu Ile Tyr Ala Phe Pro	
55 60 65	
GAT AAG CAG AAT GGC ATC GTT CCG GTG GTG GGC GTT GAT GGC TGG TCC	2334
Asp Lys Gln Asn Gly Ile Val Pro Val Val Gly Val Asp Gly Trp Ser	
70 75 80	
CGC ATC ATC AAT GAA AAC CAG CAG TTT GAT GGC ATG GAC TTT GAG CAG	2382
Arg Ile Ile Asn Glu Asn Gln Gln Phe Asp Gly Met Asp Phe Glu Gln	
85 90 95	
GAC AAT GAA TCC TGT ACA TGC CGG ATT TAC CGC AAG GAC CGT AAT CAT	2430
Asp Asn Glu Ser Cys Thr Cys Arg Ile Tyr Arg Lys Asp Arg Asn His	
100 105 110 115	
CCG ATC TGC GTT ACC GAA TGG ATG GAT GAA TGC CGC CGC GAA CCA TTC	2478
Pro Ile Cys Val Thr Glu Trp Met Asp Glu Cys Arg Arg Glu Pro Phe	
120 125 130	
AAA ACT CGC GAA GGC AGA GAA ATC ACG GGG CCG TGG CAG TCG CAT CCC	2526
Lys Thr Arg Glu Gly Arg Glu Ile Thr Gly Pro Trp Gln Ser His Pro	
135 140 145	
AAA CGG ATG TTA CGT CAT AAA GCC ATG ATT CAG TGT GCC CGT CTG GCC	2574
Lys Arg Met Leu Arg His Lys Ala Met Ile Gln Cys Ala Arg Leu Ala	
150 155 160	
TTC GGA TTT GCT GGT ATC TAT GAC AAG GAT GAA GCC GAG CGC ATT GTC	2622
Phe Gly Phe Ala Gly Ile Tyr Asp Lys Asp Glu Ala Glu Arg Ile Val	
165 170 175	
GAA AAT ACT GCA TAC ACT GCA GAA CGT CAG CCG GAA CGC GAC ATC ACT	2670
Glu Asn Thr Ala Tyr Thr Ala Glu Arg Gln Pro Glu Arg Asp Ile Thr	
180 185 190 195	
CCG GTT AAC GAT GAA ACC ATG CAG GAG ATT AAC ACT CTG CTG ATC GCC	2718
Pro Val Asn Asp Glu Thr Met Gln Glu Ile Asn Thr Leu Leu Ile Ala	
200 205 210	
CTG GAT AAA ACA TGG GAT GAC GAC TTA TTG CCG CTC TGT TCC CAG ATA	2766
Leu Asp Lys Thr Trp Asp Asp Asp Leu Leu Pro Leu Cys Ser Gln Ile	
215 220 225	
TTT CGC CGC GAC ATT CGT GCA TCG TCA GAA CTG ACA CAG GCC GAA GCA	2814
Phe Arg Arg Asp Ile Arg Ala Ser Ser Glu Leu Thr Gln Ala Glu Ala	
230 235 240	

GTA AAA GCT CTT GGA TTC CTG AAA CAG AAA GCC GCA GAG CAG AAG GTG Val Lys Ala Leu Gly Phe Leu Lys Gln Lys Ala Ala Glu Gln Lys Val 245 250 255	2852
GCA GCA TAG ATCTCGAGAA GCTTCCTGCT GAACATCAAA GGCAAGAAAA Ala Ala * 260	2911
CATCTGTTGT CAAAGACAGC ATCCTTGAAC AAGGACAATT AACAGTTAAC AAATAAAAAAC	2971
GCAAAAGAAA ATGCCGATAT CCTATTGGCA TTTTCTTTTA TTTCTTATCA ACATAAAGGT	3031
GAATCCCAIA CCTCGAGCTT CACGCTGCCG CAAGCACTCA GGGCGCAAGG GCTGCTAAAA	3091
GGAAGCGGAA CACGTAGAAA CCCAGTCCGC AGAAACGGTG CTGACCCCGG ATGAATGTCA	3151
GCTACTGGGC TATCTGGACA AGGGAAAAAC CAAGCGCAAA GAGAAAGCAG GTAGCTTGCA	3211
GTGGGCTTAC ATGGCGATAG CTAGACTGGG CGGTTTTATG GACAGCAAGC GAACCGGAAT	3271
TGCCAGCTGG GGCGCCCTCT GGTAAGGTTG GGAAGCCCTG CAAAGTAAAC TGGATGGCTT	3331
TCTTGCCGCC AAGGATCTGA TGGCGCAGGG GATCAAGATC TGATCAAGAG ACAGGATGAG	3391
GATCGTTTTG C ATG GAT ATT AAT ACT GAA ACT GAG ATC AAG CAA AAG CAT Met Asp Ile Asn Thr Glu Thr Glu Ile Lys Gln Lys His 1 5 10	3441
TCA CTA ACC CCC TTT CCT GTT TTC CTA ATC AGC CCG GCA TTT CGC GGG Ser Leu Thr Pro Phe Pro Val Phe Leu Ile Ser Pro Ala Phe Arg Gly 15 20 25	3489
CGA TAT TTT CAC AGC TAT TTC AGG AGT TCA GCC ATG AAC GCT TAT TAC Arg Tyr Phe His Ser Tyr Phe Arg Ser Ser Ala Met Asn Ala Tyr Tyr 30 35 40 45	3537
ATT CAG GAT CGT CTT GAG GCT CAG AGC TGG GCG CGT CAC TAC CAG CAG Ile Gln Asp Arg Leu Glu Ala Gln Ser Trp Ala Arg His Tyr Gln Gln 50 55 60	3585
CTC GCC CGT GAA GAG AAA GAG GCA GAA CTG GCA GAC GAC ATG GAA AAA Leu Ala Arg Glu Glu Lys Glu Ala Glu Leu Ala Asp Asp Met Glu Lys 65 70 75	3633
GGC CTG CCC CAG CAC CTG TTT GAA TCG CTA TGC ATC GAT CAT TTG CAA Gly Leu Pro Gln His Leu Phe Glu Ser Leu Cys Ile Asp His Leu Gln 80 85 90	3681
CGC CAC GGG GCC AGC AAA AAA TCC ATT ACC CGT GCG TTT GAT GAC GAT Arg His Gly Ala Ser Lys Lys Ser Ile Thr Arg Ala Phe Asp Asp Asp 95 100 105	3729
GTT GAG TTT CAG GAG CGC ATG GCA GAA CAC ATC CGG TAC ATG GTT GAA Val Glu Phe Gln Glu Arg Met Ala Glu His Ile Arg Tyr Met Val Glu 110 115 120 125	3777
ACC ATT GCT CAC CAC CAG GTT GAT ATT GAT TCA GAG GTA TAA Thr Ile Ala His His Gln Val Asp Ile Asp Ser Glu Val * 130 135	3819
AACGAGTAGA AGCTTGGCTG TTTTGGCGGA TGAGAGAAGA TTTTCAGCCT GATACAGATT	3879
AAATCAGAAC GCAGAAGCGG TGTGATAAAA CAGAATTTGC CTGGCGGCAG TAGCGCGGTG	3939
GTCCACCTG ACCCCATGCC GAACTCAGAA GTGAAACGCC GTAGCGCCGA TGGTAGTGTG	3999
GGGTCTCCCC ATGCGAGAGT AGGGAAGTGC CAGGCATCAA ATAAAACGAA AGGCTCAGTC	4059

GAAAGACTGG GCCTTTCGTT TTATCTGTTG TTTGTCCGGTG AACGCTCTCC TGAGTAGGAC 4119  
AAATCCGCCG GGAGCGGATT TGAACGTTGC GAAGCAACGG CCCGGAGGGT GGCGGGCAGG 4179  
ACGCCCCGCA TAAACTGCCA GGCATCAAAT TAAGCAGAAG GCCATCCTGA CGGATGGCCT 4239  
TTTTGCGTTT CTACAAACTC TTTTGTITAT TTTTCTAAAT ACATTCAAAT ATGTATCCGC 4299  
TCATGAGACA ATAACCCTGA TAAATGCTTC AATAATATTG AAAAAGGAAG AGTATGAGTA 4359  
TTCAACATTT CCGTGTCCGC CTTATTCCTT TTTTTCGGC ATTTTGCCTT CCTGTTTTTG 4419  
CTCACCAGAG AACGCTGGTG AAAGTAAAG ATGCTGAAGA TCAGTTGGGT GCACGAGTGG 4479  
GTTACATCGA ACTGGATCTC AACAGCGGTA AGATCCTTGA GACTTTTCGC CCCGAAGAAC 4539  
GTTTTCCAAT GATGAGCACT TTAAAGTTC TGCTATGTGG CGCGGTATTA TCCCGTGTG 4599  
ACGCCCCGCA AGAGCAACTC GGTCCGCCA TACACTATTC TCAGAATGAC TTGGTTGAGT 4659  
ACTCACCAGT CACAGAAAAG CATCTTACGG ATGGCATGAC AGTAAGAGAA TTATGCAGTG 4719  
CTGCCATAAC CATGAGTGAT AACACTGCGG CCAACTTACT TCTGACAACG ATCGGAGGAC 4779  
CGAAGGAGCT AACCGCTTTT TTGCACAACA TGGGGGATCA TGTAACTCGC CTTGATCGTT 4839  
GGGAACCGGA GCTGAATGAA GCCATACCAA ACGACGAGCG TGACACCACG ATGCCTGTAG 4899  
CAATGGCAAC AACGTTGCGC AAATATTAA CTGGCGAACT ACTTACTCTA GCTTCCCGGC 4959  
AACAAATTAAT AGACTGGATG GAGGCGGATA AAGTTGCAGG ACCACTTCTG CGCTCGGCCC 5019  
TTCCGGCTGG CTGTTTTATT GCTGATAAAT CTGGAGCCGG TGAGCGTGGG TCTCGCGGTA 5079  
TCATTGCAGC ACTGGGGCCA GATGGTAAGC CCTCCCGTAT CGTAGTTATC TACACGACGG 5139  
GGAGTCAGGC AACTATGGAT GAACGAAATA GACAGATCGC TGAGATAGGT GCCTCACTGA 5199  
TTAAGCATTG GTAAGTGTCA GACCAAGTTT ACTCATATAT ACTTTAGATT GATTACGCG 5259  
CCCTGTAGCG GCGCATTAA GCGGCGGGT GTGGTGGTTA CGCGCAGCGT GACCGCTACA 5319  
CTTGCCAGCG CCCTAGCGCC CGCTCCTTTC GCTTCTTCC CTTCTTTCT CGCCACGTTT 5379  
GCGGGCTTTC CCCGTCAAGC TCTAAATCGG GGGCTCCCTT TAGGGTTCCG ATTTAGTGCT 5439  
TTACGGCACC TCGACCCCAA AAAACTTGAT TTGGGTGATG GTTCACGTAG TGGGCCATCG 5499  
CCCTGATAGA CGTTTTTTCG CCCTTTGACG TTGGAGTCCA CGTTCTTTAA TAGTGGACTC 5559  
TTGTTCCAAA CTGAACAAC ACTCAACCCT ATCTCGGGCT ATTCTTTTGA TTTATAAGGG 5619  
ATTTTGCCGA TTTCGGCCTA TTGGTTAAAA AATGAGCTGA TTTAACAAAA ATTTACGCG 5679  
AATTTTAACA AAATATTAC GTTTACAATT TAAAAGGATC TAGGTGAAGA TCCTTTTGA 5739  
TAATCTCATG ACCAAAATCC CTTAACGTGA GTTTTCGTTT CACTGAGCGT CAGACCCCGT 5799  
AGAAAAGATC AAAGGATCTT CTTGAGATCC TTTTTTCTG CGCGTAATCT GCTGCTTGCA 5859  
AACAAAAAAA CCACCGCTAC CAGCGGTGGT TTGTTTGCCG GATCAAGAGC TACCAACTCT 5919  
TTTTCCGAAG GTAAGTGGCT TCAGCAGAGC GCAGATACCA AATACTGTCC TTCTAGTGTA 5979  
GCCGTAGTTA GGCCACCACT TCAAGAACTC TGTAGCACCG CCTACATACC TCGCTCTGCT 6039  
AATCCTGTTA CCAGTGGCTG CTGCCAGTGG CGATAAGTCG TGTCTTACCG GGTGAGACTC 6099

AAGACGATAG TTACCGGATA AGGCGCAGCG GTCGGGCTGA ACGGGGGGTT CGTGACACACA 6159  
 GCCCAGCTTG GAGCGAACGA CCTACACCGA ACTGAGATAC CTACAGCGTG AGCTATGAGA 6219  
 AAGCGCCACG CTTCCCGAAG GGAGAAAGGC GGACAGGTAT CCGGTAAGCG GCAGGGTCGG 6279  
 AACAGGAGAG CGCACGAGGG AGCTTCCAGG GGGAAACGCC TGGTATCTTT ATAGTCCTGT 6339  
 CGGGTTTCGC CACCTCTGAC TTGAGCGTCG ATTTTGTGA TGCTCGTCAG GGGGGCGGAG 6399  
 CCTATGGAAA AACGCCAGCA ACGCGGCCTT TTTACGGTTC CTGGCCTTTT GCTGGCCTTT 6459  
 TGCTCACATG TTCTTTCCTG CGTTATCCCC TGATTCTGTG GATAACCGTA TTACCGCCTT 6519  
 TGAGTGAGCT GATACCGCTC GCCGCAGCCG AACGACCGAG CGCAGCGAGT CAGTGAGCGA 6579  
 GGAAGCGGAA GAGCGCCTGA TGCGGTATTT TCTCCTTACG CATCTGTGCG GTATTTTACA 6639  
 CCGCATAGGG TCATGGCTGC GCGCCGACAC CCGCCAACAC CCGCTGACGC GCCCTGACGG 6699  
 GCTTGTCTGC TCCCGGCATC CGCTTACAGA CAAGCTGTGA CCGTCTCCGG GAGCTGCATG 6759  
 TGTCAGAGGT TTTCACCGTC ATCACCAGAA CGCGCGAGGC AGCAAGGAGA TGGCGCCCAA 6819  
 CAGTCCCCCG GCCACGGGGC CTGCCACCAT ACCCAGCCG AAACAAGCGC TCATGAGCCC 6879  
 GAAGTGGCGA GCGCGATCTT CCCCATCGGT GATGTCGGCG ATATAGGCGC CAGCAACCGC 6939  
 ACCTGTGGCG CCGGTGATGC CGGCCACGAT GCGTCCGGCG TAGAGGATCT GCTCATGTTT 6999  
 GACAGCTTAT C 7010

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 227 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Met Thr Pro Asp Ile Ile Leu Gln Arg Thr Gly Ile Asp Val Arg Ala  
 1 5 10 15  
 Val Glu Gln Gly Asp Asp Ala Trp His Lys Leu Arg-Leu Gly Val Ile  
 20 25 30  
 Thr Ala Ser Glu Val His Asn Val Ile Ala Lys Pro Arg Ser Gly Lys  
 35 40 45  
 Lys Trp Pro Asp Met Lys Met Ser Tyr Phe His Thr Leu Leu Ala Glu  
 50 55 60  
 Val Cys Thr Gly Val Ala Pro Glu Val Asn Ala Lys Ala Leu Ala Trp  
 65 70 75 80  
 Gly Lys Gln Tyr Glu Asn Asp Ala Arg Thr Leu Phe Glu Phe Thr Ser  
 85 90 95  
 Gly Val Asn Val Thr Glu Ser Pro Ile Ile Tyr Arg Asp Glu Ser Met  
 100 105 110  
 Arg Thr Ala Cys Ser Pro Asp Gly Leu Cys Ser Asp Gly Asn Gly Leu  
 115 120 125

Glu Leu Lys Cys Pro Phe Thr Ser Arg Asp Phe Met Lys Phe Arg Leu  
 130 135 140  
 Gly Gly Phe Glu Ala Ile Lys Ser Ala Tyr Met Ala Gln Val Gln Tyr  
 145 150 155 160  
 Ser Met Trp Val Thr Arg Lys Asn Ala Trp Tyr Phe Ala Asn Tyr Asp  
 155 170 175  
 Pro Arg Met Lys Arg Glu Gly Leu His Tyr Val Val Ile Glu Arg Asp  
 180 185 190  
 Glu Lys Tyr Met Ala Ser Phe Asp Glu Ile Val Pro Glu Phe Ile Glu  
 195 200 205  
 Lys Met Asp Glu Ala Leu Ala Glu Ile Gly Phe Val Phe Gly Glu Gln  
 210 215 220  
 Trp Arg  
 225

## (2) INFORMATION FOR SEQ ID NO: 13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 262 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Met Ser Thr Ala Leu Ala Thr Leu Ala Gly Lys Leu Ala Glu Arg Val  
 1 5 10 15  
 Gly Met Asp Ser Val Asp Pro Gln Glu Leu Ile Thr Thr Leu Arg Gln  
 20 25 30  
 Thr Ala Phe Lys Gly Asp Ala Ser Asp Ala Gln Phe Ile Ala Leu Leu  
 35 40 45  
 Ile Val Ala Asn Gln Tyr Gly Leu Asn Pro Trp Thr Lys Glu Ile Tyr  
 50 55 60  
 Ala Phe Pro Asp Lys Gln Asn Gly Ile Val Pro Val Val Gly Val Asp  
 65 70 75 80  
 Gly Trp Ser Arg Ile Ile Asn Glu Asn Gln Gln Phe Asp Gly Met Asp  
 85 90 95  
 Phe Glu Gln Asp Asn Glu Ser Cys Thr Cys Arg Ile Tyr Arg Lys Asp  
 100 105 110  
 Arg Asn His Pro Ile Cys Val Thr Glu Trp Met Asp Glu Cys Arg Arg  
 115 120 125  
 Glu Pro Phe Lys Thr Arg Glu Gly Arg Glu Ile Thr Gly Pro Trp Gln  
 130 135 140  
 Ser His Pro Lys Arg Met Leu Arg His Lys Ala Met Ile Gln Cys Ala  
 145 150 155 160  
 Arg Leu Ala Phe Gly Phe Ala Gly Ile Tyr Asp Lys Asp Glu Ala Glu  
 165 170 175

Arg Ile Val Glu Asn Thr Ala Tyr Thr Ala Glu Arg Gln Pro Glu Arg  
 180 185 190  
 Asp Ile Thr Pro Val Asn Asp Glu Thr Met Gln Glu Ile Asn Thr Leu  
 195 200 205  
 Leu Ile Ala Leu Asp Lys Thr Trp Asp Asp Asp Leu Leu Pro Leu Cys  
 210 215 220  
 Ser Gln Ile Phe Arg Arg Asp Ile Arg Ala Ser Ser Glu Leu Thr Gln  
 225 230 235 240  
 Ala Glu Ala Val Lys Ala Leu Gly Phe Leu Lys Gln Lys Ala Ala Glu  
 245 250 255  
 Gln Lys Val Ala Ala \*  
 260

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 139 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Met Asp Ile Asn Thr Glu Thr Glu Ile Lys Gln Lys His Ser Leu Thr  
 1 5 10 15  
 Pro Phe Pro Val Phe Leu Ile Ser Pro Ala Phe Arg Gly Arg Tyr Phe  
 20 25 30  
 His Ser Tyr Phe Arg Ser Ser Ala Met Asn Ala Tyr Tyr Ile Gln Asp  
 35 40 45  
 Arg Leu Glu Ala Gln Ser Trp Ala Arg His Tyr Gln Gln Leu Ala Arg  
 50 55 60  
 Glu Glu Lys Glu Ala Glu Leu Ala Asp Asp Met Glu Lys Gly Leu Pro  
 65 70 75 80  
 Gln His Leu Phe Glu Ser Leu Cys Ile Asp His Leu Gln Arg His Gly  
 85 90 95  
 Ala Ser Lys Lys Ser Ile Thr Arg Ala Phe Asp Asp Asp Val Glu Phe  
 100 105 110  
 Gln Glu Arg Met Ala Glu His Ile Arg Tyr Met Val Glu Thr Ile Ala  
 115 120 125  
 His His Gln Val Asp Ile Asp Ser Glu Val \*  
 130 135



Table 1: Sequences of Oligos for PCR

## Figure 3ab

left: TGACCCCTCACAAGGAGACGACCTTCCATGACCGAGTACAAGAGGGATGTAACGCACTGA

right: TACAAATGTGGTATGGCTGATTATGATCCTCTAGAGTCGGTGCTCACTGCCCCGCTTTCCA

template: pJP5603

targeting vector: pSV-paz11

## Figure 3c

a-left: CTTCCATGACCGAGTACAAGAGGGATGTAACGCACTGA

a-right: ATGATCCTCTAGAGTCGGTGCTCACTGCCCCGCTTTCCA

b-left: AGACGACCTTCCATGACCGAGTACAAGAGGGATGTAACGCACTGA

b-right: GCTGATTATGATCCTCTAGAGTCGGTGCTCACTGCCCCGCTTTCCA

c-left: CACAAGGAGACGACCTTCCATGACCGAGTACAAGAGGGATGTAACGCACTGA

c-right: TGGTATGGCTGATTATGATCCTCTAGAGTCGGTGCTCACTGCCCCGCTTTCCA

d-left: TGACCCCTCACAAGGAGACGACCTTCCATGACCGAGTACAAGAGGGATGTAACGCACTGA

d-right: TACAAATGTGGTATGGCTGATTATGATCCTCTAGAGTCGGTGCTCACTGCCCCGCTTTCCA

e-left:

CACGCCCCTGACCCCTCACAAGGAGACGACCTTCCATGACCGAGTACAAGAGGGATGTAACGCACTGA

e-right:

TAAACCTCTACAAATGTGGTATGGCTGATTATGATCCTCTAGAGTCGGTGCTCACTGCCCCGCTTTCCA

f-left:

TCCCCTGACCCACGCCCCTGACCCCTCACAAGGAGACGACCTTCCATGACCGAGTACAAGAGGGATGT  
AACGCACTGA

f-right:

TAAAGCAAGTAAACCTCTACAAATGTGGTATGGCTGATTATGATCCTCTAGAGTCGGTGCTCACTGCC  
CGCTTTCCA

template: pJP5603

targeting vector: pSV-paz11

## Figure 3d

a-left:

TCATCCTCTGCATGGTCAGGTCATGGATGAGCAGACGATGGTGCAAGGCTGCTAAAGGAA

a-right:

TAATGCGAACAGCGCACGGCGTTAAAGTTGTTCTGCTTCATCAGCAGGATGGCGAAGAACTCCAGCAT

b-left:

CACGAGCATCATCCTCTGCATGGTCAGGTCATGGATGAGCAGACGATGGCAAGGGCTGCTAAAGGAA

b-right:

TAATGCGAACAGCGCACGGCGTTAAAGTTGTTCTGCTTCATCAGCAGGATGGCGAAGAACTCCAGCAT

c-left:

TTAACCGTCACGAGCATCATCCTCTGCATGGTCAGGTCATGGATGAGCACAAGGGCTGCTAAAGGAA

c-right:

TAATGCGAACAGCGCACGGCGTTAAAGTTGTTCTGCTTCATCAGCAGGATGGCGAAGAACTCCAGCAT

d-left:

TGCTGCTGAACGGCAAGCCGTTGCTGATTGAGGCGTTAACCGTCACGACAAGGGCTGCTAAAGGAA

d-right:

TAATGCGAACAGCGCACGGCGTTAAAGTTGTTCTGCTTCATCAGCAGGATGGCGAAGAACTCCAGCAT

e-left:

TCTCTATCGTGCGGTGGTTGAACTGCACACCGCCGACGGCAGCTGATTCAAGGGCTGCTAAAGGAA

e-right:

TAATGCGAACAGCGCACGGCGTTAAAGTTGTTCTGCTTCATCAGCAGGATGGCGAAGAACTCCAGCAT

f-left:

TGGAGTGACGGCAGTTATCTGGAAGATCAGGATAATGTTGGCGGATGAGCGCAAGGGCTGCTAAAGGAA

f-right:

TAATGCGAACAGCGCACGGCGTTAAAGTTGTTCTGCTTCATCAGCAGGATGGCGAAGAACTCCAGCAT

g-left:

TGCATTCTAGTTGTGGTTTGTCCAAACTCATCAATGTATCTTATCATGTCAAGGGCTGCTAAAGGAA

g-right:

TAATGCGAACAGCGCACGGCGTTAAAGTTGTTCTGCTTCATCAGCAGGATGGCGAAGAACTCCAGCAT

h-left:

TGCATTCTAGTTGTGGTTTGTCCAAACTCATCAATGTATCTTATCATGTCAAGGGCTGCTAAAGGAA

h-right:

TATTTTGTACACCAGACCAACTGGTAATGGTAGCGACCGGCGCTCAGCTGGCGAAGAACTCCAGCAT

template: pJP5603

targeting vector: pSV-paz11

Figure 4

left:

TCAAGTGAGAAATCACCATGAGTGACGACTGAATCCGGTGAGAATGGCAAAAGCTTATGCCCACCAGC  
TGGTATGGCTGATTATGATC

right:

TCCAACATGGATGCTGATTTATATGGGTATAAATGGGCTCGCGATAATGTCGGGCAATCAGGTGCGACA  
ATCTACCACCAGCTCTTTTCTACGGGGTCTGACGC

template: pBR322

targeting vector: Hoxa-PI

Figure 5

left:

TGCAGCTGGCACGACAGGTTTCCCGACTGGAAAGCGGGCAGTTAATACGACTCACTATAGGGAGAACA  
GGAAACAGCTATGCCCATAACACCCAGAGTA

right:

TGC CGCCTACAGGGCGCGTCCATTCCGCCATTCAGGCCTGACTCACTAGTGATGGTGATGGTGATGTGG  
GGGGTGGCGCTCAGT

template: pmtrx (a pBluescript vector carrying mouse trithorax cDNA)

targeting vector: pZero2.1

Figure 6

left:

TGAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGGAGAAAAAATCACT  
GGATATACCACCG

right:

TACAGGGCGCGTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACGCCCCGCCCTGC  
CACTCATCGCA

template: pMAK705

targeting vector: pBAD-24 backbone Amp resistant gene

Figure 8

i:

TGCCAAGCTTGACCCACTGTGGAAGTGTTCCAAAAAGCGGGAAGGCTCTTGAGCTACTTCACTAACAAC  
CGG

g:

TCACCATCTTCGGGCCATTTGTAGACTGGAATATTTGAGCTATGAGTGTGCTACTTCACTAACAACCG  
G

h:

TGGCCCCAGGGTGACGCGGACATGGAGTTGTGCGCCAGGGCACTGGTCCATGAGAGTGCCAAGCTACTC  
GCGAC

template: pKaZ

targeting vector: Hoxa-PI

Figure 9

j:

TAATAGCGAAGAGGCCCGCACCGATCGCCCTTCCCAACAGTTGCGCAGCCTGAATGGCGAATGGCGCT  
TTGCTGGTTTATAACTTCGTATAGCATACATTATACGAAGTTATGGGCTGCTAAAGGAAGCGGAACAC  
G

k:

TGGCAGTTCAGGCCAATCCGCGCCGGATGCGGTGTATCGCTCGCCACTTCAACATCAACGGTAATCGCC  
ATTTGACCATATAACTTCGTATAATGTATGCTATACGAAGTTATCCCCAGAGTCCCGCTCAGAAGAACT

template: pJP5603

targeting vector: JC9604 chromosome

Figure 10

l:

TAGCTTGGCACTGGCCGTCGTTTTACAACGTCGTGACTGGGAAAACCCTGGCGTTACCCAACCCATCAC  
ATATACCTGCCGTTCACTAT

m:

TATCGGTGGCCGTGGTGTGCGCTCCGCCGCCTTCATACTGCACCGGGCGGGAAGGCGATTCCGAAGCCC  
AACCTTTCATAGAAGCC

template: pIB279

targeting vector: pSV-paX1

l\*: GCTTGGCACTGGCCGTCGTTTTACAACGTCGTGACTGGGAA

m\*:

TCGGTGGCCGTGGTGTGCGCTCCGCCGCCTTCATACTGCACCGGGCGGGAAGGATCCACAGATTTGATC  
CAGCGATACAGC

template: pSV-paz11

targeting vector: pSV-sacB-neo

Figure 11

n:

TACCGCATTAAAGCTTATCGATGATAAGCTGTCAAACATGAGAATTGACCCGGAACCCTTCTCGAGGAA  
GTTCTATTCTCTAGAAAGTATAGGAACCTCCGAATAAATACCTGTGACGGAAGATCACTT

p:

TTCCCTCAAGAATTTTACTCTGTCAGAAACGGCCTTAACGACGTAGTCGAGGGACCTAGAAGTTCCTAT  
ACTTTCTAGAGAATAGGAACCTCATTATCACTTATTCAGGCGTAGCACCAGGCG

template: pMAK705

targeting vector: Hoxa-P1

Figure 12

left:

TGAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGGAGAAAAAATCACT  
GGATATACCACCG

right:

TACAGGGCGCGTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACGCCCCGCCCTGC  
CACTCATCGCA

template: pMAK705

targeting vector: pBAD-24 backbone Amp resistant gene

**THIS PAGE BLANK (USPTO)**




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : <b>C12N 15/10, 15/90</b>		(11) International Publication Number: <b>WO 99/29837</b>
<b>A3</b>		(43) International Publication Date: 17 June 1999 (17.06.99)
(21) International Application Number: PCT/EP98/07945		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 7 December 1998 (07.12.98)		
(30) Priority Data: 97121462.2 5 December 1997 (05.12.97) EP 98118756.0 5 October 1998 (05.10.98) EP		
(71) Applicant (for all designated States except US): EUROPAISCHES LABORATORIUM FÜR MOLEKULARBIOLOGIE (EMBL) [DE/DE]; Meyerhofstrasse 1, D-69117 Heidelberg (DE).		
(72) Inventors; and (75) Inventors/Applicants (for US only): STEWART, Francis [AU/DE]; Lärchenweg 3, D-69181 Leimen (DE). ZHANG, Youming [CN/DE]; Friedrich-Ebert-Anlage 51c, D-69117 Heidelberg (DE). BUCHHOLZ, Frank [DE/DE]; Neuenkirchener Weg 44a, D-28779 Bremen (DE).		
(74) Agents: WEICKMANN, H. et al.; Kopernikusstrasse 9, D-81679 München (DE).		<p><b>Published</b></p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> <p>(88) Date of publication of the international search report: 22 July 1999 (22.07.99)</p>

(54) Title: NOVEL DNA CLONING METHOD RELYING ON THE E. COLI REC<sup>E</sup>/REC<sup>T</sup> RECOMBINATION SYSTEM

## (57) Abstract

The invention refers to a novel method for cloning DNA molecules using a homologous recombination mechanism between at least two DNA molecules comprising: a) providing a host cell capable of performing homologous recombination, b) contacting in said host cell a first DNA molecule which is capable of being replicated in said host cell with a second DNA molecule comprising at least two regions of sequence homology to regions on the first DNA molecule, under conditions which favour homologous recombination between said first and second DNA molecules and c) selecting a host cell in which homologous recombination between said first and second DNA molecules has occurred. In particular, it relies on the use of the E. coli RecE and RecT proteins, the bacteriophage Red-alpha and Red-beta proteins, or the phage P22 recombination system. The beneficial effects of concomitant expression of the RecBC inhibitor genes (e.g. Red-Gamma) is also exemplified.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

Original Application No

PCT/EP 98/07945

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/10 C12N15/90

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched: classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	OLINER J.D. ET AL.: "In vivo Cloning of PCR Products in E. coli" NUCLEIC ACIDS RESEARCH, vol. 21, no. 22, 1993, pages 5192-5197, XP002064297 cited in the application	1-6, 8-13, 19, 20, 22-27, 34, 35, 37-42, 46-48, 50
Y	See page 5192, column 2, line 8 - line 31, and discussion section	18, 28
Y	NUSSBAUM A ET AL: "Restriction-stimulated homologous recombination of plasmids by the RecE pathway of Escherichia coli" GENETICS, vol. 130, no. 1, January 1992, pages 37-49, XP002103370 see figures 1, 7	18, 28

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

### Special categories of cited documents

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"S" document member of the same patent family

Date of the actual completion of the international search

21 May 1999

Date of making of the international search report

04/06/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Lonnoy, O

## C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No. *
P, X	ZHANG Y ET AL: "A new logic for DNA engineering using recombination in Escherichia coli" NATURE GENETICS, vol. 20, no. 2, October 1998, pages 123-128. XP002103371 see the whole document ---	1-50
A	KOLODNER, R ET AL: "Homologous pairing proteins encoded by the Escherichia coli recE and recT genes" MOLECULAR MICROBIOLOGY, vol. 11, no. 1, 1994, pages 23-30, XP002064301 cited in the application ---	
A	LUISI-DELUCA C ET AL: "Genetic and physical analysis of plasmid recombination in recB recC sbcB and recB recC sbcA Escherichia coli K-12 mutants" GENETICS, vol. 122, 1989, pages 269-278. XP002064305 ---	
A	DEGRYSE E: "Evaluation of Escherichia coli recBC sbcBC mutants for cloning by recombination in vivo" JOURNAL OF BIOTECHNOLOGY, vol. 2, no. 39, 15 April 1995, page 181-187 XP004036984 ---	
A	YANG X ET AL: "Homologous recombination based modification in Escherichia coli and germ line transmission in transgenic mice of a bacterial artificial chromosome" NATURE BIOTECHNOLOGY, vol. 15, September 1997, pages 859-865, XP002103372 ---	
A	MURPHY K: "Lambda Gam protein inhibits the helicase and Chi-stimulated recombination activities of Escherichia coli RecBCD enzyme" JOURNAL OF BACTERIOLOGY, vol. 173, no. 18, September 1991, pages 5808-5821. XP002103373 -----	






## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/10, 15/90</b>		<b>A3</b>	(11) International Publication Number: <b>WO 99/29837</b>
			(43) International Publication Date: 17 June 1999 (17.06.99)
(21) International Application Number: PCT EP98/07945		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 7 December 1998 (07.12.98)		Published <i>With international search report</i> <i>With amended claims.</i>	
(30) Priority Data: 97121462.2 5 December 1997 (05.12.97) EP 98118756.0 5 October 1998 (05.10.98) EP		(88) Date of publication of the international search report: 22 July 1999 (22.07.99)	
(71) Applicant (for all designated States except US): EUROPAISCHES LABORATORIUM FÜR MOLEKULARBIOLOGIE (EMBL) [DE/DE]; Meyerhofstrasse 1, D-69117 Heidelberg (DE).		Date of publication of the amended claims: 16 September 1999 (16.09.99)	
(72) Inventors; and (75) Inventors/Applicants (for US only): STEWART, Francis [AU/DE]; Lärchenweg 3, D-69181 Leimen (DE). ZHANG, Youming [CN/DE]; Friedrich-Ebert-Anlage 51e, D-69117 Heidelberg (DE). BUCHHOLZ, Frank [DE/DE]; Neuenkirchener Weg 44a, D-28779 Bremen (DE).			
(74) Agents: WEICKMANN, H. et al.; Kopernikusstrasse 9, D-81679 München (DE).			

(54) Title: NOVEL DNA CLONING METHOD RELYING ON THE E. COLI REC/RECT RECOMBINATION SYSTEM

## (57) Abstract

The invention refers to a novel method for cloning DNA molecules using a homologous recombination mechanism between at least two DNA molecules comprising: a) providing a host cell capable of performing homologous recombination, b) contacting in said host cell a first DNA molecule which is capable of being replicated in said host cell with a second DNA molecule comprising at least two regions of sequence homology to regions on the first DNA molecule, under conditions which favour homologous recombination between said first and second DNA molecules and c) selecting a host cell in which homologous recombination between said first and second DNA molecules has occurred. In particular, it relies on the use of the E. coli RecE and RecT proteins, the bacteriophage Red-alpha and Red-beta proteins, or the phage P22 recombination system. The beneficial effects of concomitant expression of the RecBC inhibitor genes (e.g. Red-Gamma) is also exemplified.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## AMENDED CLAIMS

[received by the International Bureau on 04 August 1999 (04.08.99);  
original claims 1-50 replaced by new claims 1-64 (10 pages)]

1. A method for cloning DNA molecules in procaryotic cells comprising the steps of:
  - a) providing a procaryotic host cell capable of performing homologous recombination,
  - b) contacting in said host cell a circular first DNA molecule which is capable of being replicated in said host cell with a second DNA molecule comprising at least two regions of sequence homology to regions on the first DNA molecule, under conditions which favour homologous recombination between said first and second DNA molecules and
  - c) selecting a host cell in which homologous recombination between said first and second DNA molecules has occurred.
2. The method according to claim 1 wherein the homologous recombination occurs via the recET cloning mechanism.
3. The method according to claim 2 wherein the host cell is capable of expressing recE and recT genes.
4. The method according to claim 3 wherein the recE and recT genes are selected from E.coli recE and recT genes or from  $\lambda$  red $\alpha$  and red $\beta$  genes.
5. The method according to claim 3 or 4 wherein the host cell is transformed with at least one vector capable of expressing recE and/or recT genes.
6. The method of claim 3, 4 or 5 wherein the expression of the recE and/or recT genes is under control of a regulatable promoter.

7. The method of claim 5 or 6 wherein the *recT* gene is overexpressed versus the *recE* gene.
8. The method according to any one of claims 3 to 7 wherein the *recE* gene is selected from a nucleic acid molecule comprising
  - (a) the nucleic acid sequence from position 1320 (ATG) to 2159 (GAC) as depicted in Fig.7B,
  - (b) the nucleic acid sequence from position 1320 (ATG) to 1998 (CGA) as depicted in Fig.13B,
  - (c) a nucleic acid encoding the same polypeptide within the degeneracy of the genetic code and/or
  - (d) a nucleic acid sequence which hybridizes under stringent conditions with the nucleic acid sequence from (a), (b) and/or (c).
9. The method according to any one of claims 3 to 8 wherein the *recT* gene is selected from a nucleic acid molecule comprising
  - (a) the nucleic acid sequence from position 2155 (ATG) to 2961 (GAA) as depicted in Fig.7B,
  - (b) the nucleic acid sequence from position 2086 (ATG) to 2868 (GCA) as depicted in Fig.13B,
  - (c) a nucleic acid encoding the same polypeptide within the degeneracy of the genetic code and/or
  - (d) a nucleic acid sequence which hybridizes under stringent conditions with the nucleic acid sequences from (a), (b) and/or (c).
10. The method according to any one of the previous claims wherein the host cell is a gram-negative bacterial cell.
11. The method according to claim 10 wherein the host cell is an *Escherichia coli* cell.

12. The method according to claim 11 wherein the host cell is an *Escherichia coli* K12 strain.
13. The method according to claim 12 wherein the *E. coli* strain is selected from JC 8679 and JC 9604.
14. The method according to any one of the previous claims wherein the host cell further is capable of expressing a *recBC* inhibitor gene.
15. The method according to claim 14 wherein the host cell is transformed with a vector expressing the *recBC* inhibitor gene.
16. The method according to claim 14 or 15 wherein the *recBC* inhibitor gene is selected from a nucleic acid molecule comprising
  - (a) the nucleic acid sequence from position 3588 (ATG) to 4002 (GTA) as depicted in Fig.13B,
  - (b) a nucleic acid encoding the same polypeptide within the degeneracy of the genetic code and/or
  - (c) a nucleic acid sequence which hybridizes under stringent conditions (as defined above) with the nucleic acid sequence from (a) and/ or (b).
17. The method according to any one of claims 13 to 16 wherein the host cell is a prokaryotic *recBC* + cell.
18. The method according to any one of the previous claims wherein the first DNA molecule is an extrachromosomal DNA molecule containing an origin of replication which is operative in the host cell.
19. The method according to claim 18 wherein the first DNA molecule is selected from plasmids, cosmids, P1 vectors, BAC vectors and PAC vectors.

20. The method according to any one of claims 1-18 wherein the first DNA molecule is a host cell chromosome.
21. The method according to any one of the previous claims wherein the second DNA molecule is linear.
22. The method according to any one of the previous claims wherein the regions of sequence homology are at least 15 nucleotides each.
23. The method according to one of claims 1 to 16 wherein the second DNA molecule is obtained by an amplification reaction.
24. The method according to one of the previous claims wherein the first and/or second DNA molecules are introduced into the host cells by transformation.
25. The method according to claim 24 wherein the transformation method is electroporation.
26. The method according to one of claims 1 to 25 wherein the first and second DNA molecules are introduced into the host cell simultaneously by co-transformation.
27. The method according to one of claims 1 to 25 wherein the second DNA molecule is introduced into a host cell in which the first DNA molecule is already present.
28. The method according to one of the previous claims wherein the second DNA molecule contains at least one marker gene placed between the two regions of sequence homology and wherein homologous recombination is detected by expression of said marker gene.

**AMENDED SHEET (ARTICLE 19)**

29. The method according to claim 28 wherein gene presence is selected from antibiotic resistance genes, deficiency complementation genes and reporter genes.
30. The method of any one of claims 1 to 29 wherein the first DNA molecule contains at least one marker gene between the two regions of sequence homology and wherein homologous recombination is detected by lack of expression of said marker gene.
31. The method of any one of claims 1 to 30 wherein said marker gene is selected from genes which, under selected conditions, convey a toxic or bacteriostatic effect on the cell, and reporter genes.
32. A method according to any one of the previous claims wherein the first DNA molecule contains at least one target site for a site specific recombinase between the two regions of sequence homology and wherein homologous recombination is detected by removal of said target site.
33. A method for cloning DNA molecules comprising the steps of:
  - (a) providing a source of RecE and RecT proteins,
  - (b) contacting a first DNA molecule which is capable of being replicated in a suitable host cell with a second DNA molecule comprising at least two regions of sequence homology to regions on the first DNA molecule, under conditions which favour homologous recombination between said first and second DNA molecules and
  - (c) selecting DNA molecules in which homologous recombination between said first and second DNA molecules has occurred.
34. The method of claim 33 wherein said RecE and RecT or proteins are selected from E.coli RecE and RecT proteins or from phage  $\lambda$  Red $\alpha$  and Red $\beta$  proteins.

35. The method of claim 33 or 34 wherein the recombination occurs in vitro.
36. The method of claim 33 or 34 wherein the recombination occurs in vivo.
37. A method for making a recombinant DNA molecule comprising introducing into a prokaryotic host cell a circular first DNA molecule which is capable of being replicated in said host cell, and introducing a second DNA molecule comprising a first and a second region of sequence homology to a third and fourth region, respectively, on the first DNA molecule, said host cell being capable of performing homologous recombination, such that a recombinant DNA molecule is made, said recombinant DNA molecule comprising the first DNA molecule wherein the sequences between said third and fourth regions have been replaced by sequences between the first and second regions of the second DNA molecule.
38. The method according to claim 37 which further comprises detecting the recombinant DNA molecule.
39. A method for making a recombinant DNA molecule comprising introducing into a prokaryotic host cell, containing a chromosomal first DNA molecule, a second DNA molecule comprising a first and a second region of sequence homology to a third and a fourth region, respectively, on the host chromosomal first DNA molecule, said host cell being capable of performing homologous recombination, such that a recombinant DNA molecule is made, said recombinant DNA molecule comprising the chromosomal first DNA molecule wherein the sequences between said third and fourth regions have been replaced by sequences between the first and second regions of the second DNA molecule.



40. The method according to claim 39 which further comprises detecting the recombinant DNA molecule.
41. The method according to any one of claims 37 to 40, wherein the host cell is capable of expressing RecE and RecT proteins or  $\lambda$ exo and  $\lambda\beta$  proteins.
42. A method for cloning DNA molecules comprising the steps of:
- (a) contacting in vitro a first DNA molecule with a second DNA molecule comprising at least two regions of sequence homology to regions on the first DNA molecule, in the presence of RecE and RecT proteins and under conditions which favour homologous recombination between said first and second DNA molecules; and
  - (b) selecting a DNA molecule in which homologous recombination between said first and second DNA molecules has occurred.
43. A method for making a recombinant DNA molecule comprising contacting in vitro a first DNA molecule with a second DNA molecule comprising a first and a second region of sequence homology to a third and a fourth region on the first DNA molecule, in the presence of RecE and RecT proteins and under conditions in which homologous recombination can occur, such that a recombinant DNA molecule is made, said recombinant DNA molecule comprising the first DNA molecule wherein the sequences between said third and fourth regions have been replaced by sequences between the first and second regions of the second DNA molecule.
44. The method of claim 42, which further comprises between steps (a) and (b) the step of introducing the product step (a) into a cell, wherein recombination occurs in the cell.

**THIS PAGE BLANK (USPTO)**